

A population of high-redshift type 2 quasars – I. Selection criteria and optical spectra

Alejo Martínez-Sansigre,^{1★} Steve Rawlings,¹ Mark Lacy,² Dario Fadda,²
Matt J. Jarvis,¹ Francine R. Marleau,² Chris Simpson^{3,4} and Chris J. Willott⁵

¹*Astrophysics, Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH*

²*Spitzer Science Center, California Institute of Technology, MS220-6, 1200 E. California Boulevard, Pasadena, CA 91125, USA*

³*Department of Physics, University of Durham, South Road, Durham DH1 3LE*

⁴*Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD*

⁵*Herzberg Institute of Astrophysics, National Research Council, 5071 West Saanich Rd, Victoria, B.C. V9E 2E7, Canada*

Accepted 2006 May 15. Received 2006 May 15; in original form 2006 January 20

ABSTRACT

We discuss the relative merits of mid-infrared and X-ray selection of type 2 quasars. We describe the mid-infrared, near-infrared and radio selection criteria used to find a population of redshift $z \sim 2$ type 2 quasars which we previously argued suggests that most supermassive black hole growth in the Universe is obscured. We present the optical spectra obtained from the William Herschel Telescope, and we compare the narrow emission-line luminosity, radio luminosity and maximum size of jets to those of objects from radio-selected samples. This analysis suggests that these are genuine radio-quiet type 2 quasars, albeit the radio-bright end of this population. We also discuss the possibility of two different types of quasar obscuration, which could explain how the ~ 2 – 3 :1 ratio of type 2 to type 1 quasars preferred by modelling our population can be reconciled with the ~ 1 :1 ratio predicted by unified schemes.

Key words: galaxies: active – galaxies: nuclei – quasars: general.

1 INTRODUCTION

For a long time, a population of obscured (type 2) radio-quiet quasars has been postulated from modelling the hard (energies ≥ 1 keV) X-ray background as the sum of the emission from extragalactic sources. Comastri et al. (1995) assumed a population of absorbed active galactic nuclei (AGN; in this paper we use the term to encompass both Seyferts and quasars), with the same intrinsic spectrum and evolution as unobscured AGN, but with a range of obscuring hydrogen column densities, N_{H} , between 10^{25} and 10^{29} m⁻². The resulting best-fitting ratio for absorbed ($N_{\text{H}} > 10^{26.5}$ m⁻²) to unabsorbed ($N_{\text{H}} < 10^{26.5}$ m⁻²) AGN was found to be ~ 3 :1. The model assumed the same distribution of obscuring column densities and redshifts for the low X-ray luminosity objects ($L_{\text{X}} < 10^{37}$ W; the Seyferts) and the high X-ray luminosity objects ($L_{\text{X}} > 10^{37}$ W; the quasars), including a population of highly obscured AGN (the Seyfert 2s and type 2 quasars).

According to unified schemes (Antonucci 1993), type 2 quasars would consist of quasars with the symmetry axes perpendicular to the observer's line of sight, so that the dusty torus around the accretion disc would be viewed edge on, obscuring the optically bright accretion disc. The gas present in this torus will obscure the X-rays via photoelectric absorption and Compton scattering. This

unfavourable orientation means type 2 quasars do not outshine their host galaxy at (rest-frame) ultraviolet (UV) or optical wavelengths, as is the case with the unobscured (type 1) quasars. Type 2 quasars are therefore indistinguishable from normal galaxies in optical imaging surveys. If the torus has a half-opening angle of $\sim 40^\circ$, as appears to be the case in radio-loud samples (e.g. Willott et al. 2000), the population of type 2s is expected to be comparable in size to the type 1 population and not to outnumber it by a factor of ~ 3 .

A modification to the standard unified scheme is the ‘receding torus’ model (Lawrence 1991) in which the more luminous AGN sublimate the dust in the inner edge of the torus out to larger distances than the less luminous AGN. This leads to a larger opening angle for more luminous AGN, so the fraction of type 1 quasars increases as a function of bolometric luminosity. Such luminosity dependence of the type 2 to type 1 ratio is indeed found, for example, in radio-selected samples by Willott et al. (2000), in X-ray-selected samples by Ueda et al. (2003), and in spectroscopically selected samples by Simpson (2005).

More recent modelling of the hard X-ray background has consistently shown signs of a large population of obscured AGN (Wilman & Fabian 1999; Worsley et al. 2005) and synthesis models have generally required a type 2 to type 1 ratio ~ 3 – 4 :1 (Gilli, Salvati & Hasinger 2001; Ueda et al. 2003; Treister & Urry 2005). However, the bulk of the sources required to fit the unresolved X-ray background has moderate redshifts ($z = 0.5$ – 1) and X-ray luminosities ($L_{\text{X}} \sim 10^{36}$ W) characteristic of Seyferts rather than

★E-mail: a.martinez-sansigre1@physics.oxford.ac.uk

quasars. The latest studies of the hard X-ray background indeed suggest a luminosity-dependent Compton-thin type 2 to type 1 ratio, which decreases to $\lesssim 1:1$ at the higher luminosities corresponding to quasars (Ueda et al. 2003; Treister & Urry 2005). The true type 2 to type 1 ratio will remain unknown until the Compton-thick population is unveiled.

Deep hard X-ray surveys have now revealed a large population of obscured AGN (e.g. Alexander et al. 2003), but again most of these sources have moderate X-ray luminosities ($L_X < 10^{37}$ W) and are better described as Seyfert 2s rather than type 2 quasars. Amongst objects with X-ray luminosities large enough to qualify as quasars, the ratio of type 2 to type 1 objects appears to be only $\sim 1:1$ (e.g. Zheng et al. 2004), although redshift completeness at faint optical magnitudes ($m_R > 23$, Vega) may still be an issue. Despite the great progress with the deep hard X-ray surveys, at high energies (above 6–8 keV) only ~ 50 per cent of the X-ray background is accounted for by individual sources, suggesting a substantial population of Compton-thick AGN which contribute to the hard X-ray background but are missing from current flux-limited X-ray surveys (Worsley et al. 2005). Indeed, a good example is the type 2 quasar IRAS FSC 10214+4724: this object, found in an IRAS 60- μ m survey (Rowan-Robinson et al. 1991), has recently been found to be Compton thick (Alexander et al. 2005).

An alternative strategy for looking for type 2 objects is to look for the mid-infrared emission. IRAS FSC 10214+4724, at $z \sim 2.3$, is a classic type 2 object (e.g. Serjeant et al. 1998). However, due to the limited sensitivity of IRAS, this object was only detected due to a huge gravitational lens magnification (~ 50) of the flux (Broadhurst & Lehar 1995). The dramatic increase of mid-infrared sensitivity allowed by the *Spitzer* Space Telescope (Werner et al. 2004) means that similar objects can now be found without the ‘benefit’ of gravitational lensing, and mid-infrared selection should be sensitive to the Compton-thick quasars missed by X-rays.

Following the success of mid-infrared selection in finding type 2 radio-quiet quasars at $z \sim 0.5$ (Lacy et al. 2004, 2005a), we combined data from the Multi-Band Imaging Photometer for *Spitzer* (MIPS) instrument at 24 μ m (Marleau et al. 2004; Fadda et al. 2006), IRAC at 3.6 μ m (Lacy et al. 2005b) and from the Very Large Array (VLA) at 1.4 GHz (Condon et al. 2003) and devised strict selection criteria to hunt for higher redshift type 2s (Martínez-Sansigre et al. 2005). The trade-off between our method and X-ray selection is that our quasars are intrinsically more luminous than AGN found in the deep hard X-ray surveys.

The format of this paper is as follows. In Section 2, we compare X-ray and mid-infrared selection of obscured quasars. In Section 3, we explain in detail the selection criteria which we used to find our sample of type 2 quasars. Section 4 deals with the observations, while Section 5 describes the optical spectra and Section 6 compares the spectroscopic and photometric redshifts. In Section 7, we make the first composite radio-quiet type 2 spectrum. Section 8 is dedicated to comparing our sample to the well-studied radio-loud samples. Finally, Section 9 summarizes and discusses the implications of this population of type 2 quasars. Throughout this paper, we adopt a Lambda cold dark matter (Λ CDM) cosmology with the following parameters: $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1}) = 0.7$; $\Omega_m = 0.3$; $\Omega_\Lambda = 0.7$.

2 COMPARISON BETWEEN X-RAY AND MID-INFRARED SELECTION

In the unified scheme for AGN (Antonucci 1993), the central engine is surrounded by a torus of dust which will obscure the broad-line re-

gion from certain lines of sight. The optical classification for quasars is that type 1s should show broad lines (and strong optical-UV continuum), while the type 2s are those objects where the dusty torus is obscuring the broad-line region, so only narrow lines are seen. This classification is concerned with the optical/UV properties, and Simpson, Rawlings & Lacy (1999) found that the dividing extinction for radio-selected (3CR and 3CRR) objects corresponded to $A_V \gtrsim 5$, with a low fraction of objects with A_V in the range 5–15 and a large fraction with $A_V \gtrsim 15$. Note also that the Simpson et al. (1999) study found that most type 1s ($A_V \lesssim 5$) had $A_V \sim 0$, with only a small fraction lightly reddened ($A_V \sim 1$ –5). Gas and dust are responsible for obscuring the X-rays emitted from the central engine, and the geometrical distribution of the gas is almost certainly different from that of the dust. It is possible to imagine lines of sight which ‘graze’ the dusty torus, leading to small amounts of extinction in the optical, but which go through significant additional amounts of gas. Such a line of sight would lead to a quasar classified as ‘reddened’ (not type 2) in the optical ($A_V \simeq 1$ –5) but as type 2 ($N_H \gtrsim 10^{26} \text{ m}^{-2}$) in X-rays. The converse situation, where an optically obscured quasar is barely absorbed in X-rays, is harder to envisage: any line of sight that passes through significant amounts of dust will lead to significant absorption in the X-rays. It is therefore unlikely that optical type 2 quasars will have negligible X-ray absorption. However, in edge-on sources, the nuclear soft X-ray emission can plausibly be scattered from an optically thin material located above or below the plane of the torus, leading to an optically obscured AGN with a soft X-ray spectrum. Finally, obscuring material in the host galaxy of the AGN can further complicate interpretation. Independently of the orientation of the torus, dust and gas in the host galaxy can cause extinction and absorption with a huge range in A_V and N_H .

X-ray and optical definitions of obscured AGN are therefore slightly different and the range of gas-to-dust ratios found by comparing dust reddening in the optical or near-infrared and X-ray absorption suggests they are not always matched (e.g. Willott et al. 2003a; Urrutia et al. 2005; Wilkes et al. 2005). We note that we have used a mid-infrared plus radio selection, but since the role of the radio criterion was to avoid non-AGN contaminants (see Section 3), in this Section we can proceed to consider our technique as basically mid-infrared selection. The effect of adding the radio flux density cuts is to constrain the type 2 quasars selected to those at the high end of the radio-to-optical correlation for radio-quiet quasars of Cirasuolo et al. (2003), as described in Section 3 and in Martínez-Sansigre et al. (2005). It is of course plausible that radio luminosity correlates in some complicated way with joint mid-infrared, X-ray detectability but we ignore this possibility here.

This section aims to compare the ‘merits’ of mid-infrared and X-ray selection by considering a model quasar (described in Section 2.1) with varying amounts of dust extinction (Section 2.2) and different gas-to-dust ratios (Section 2.3). We also consider the effects of different types of dust (Section 2.5) and the orientation dependence of 24- μ m emission (Section 2.6).

2.1 Model quasar

We choose this quasar to have $M_B = -25.7$ since this corresponds to the break in the optical quasar luminosity function at $z = 2$ (Croom et al. 2004), the redshift of interest in the analysis in Martínez-Sansigre et al. (2005). The model quasar is assumed to have an intrinsic unreddened type 1 spectral energy distribution (SED), which we take from Rowan-Robinson (1995). This model covers the range between the (rest-frame) far-infrared and 4 keV. We need to

complement this with a SED that goes into the hard X-rays, so we assume the Madau, Ghisellini & Fabian (1994) spectrum with the form $L_\nu \propto \nu^{-0.9} \exp(-\nu/\nu_c)$ with $h\nu_c = 360$ keV. This intrinsic unabsorbed type 1 spectrum is chosen as it was used by Wilman & Fabian (1999) for the models we use for absorbed X-ray spectra. This is practically a flat X-ray SED, and in the range probed by observations at $24\ \mu\text{m}$ (for $z \leq 5$), the SED is also flat ($L_\nu \propto \nu^{-1}$ so $\nu L_\nu \propto \nu^0$). We choose the normalization to match the Rowan-Robinson (1995) SED at 2 keV.

2.2 Obscuration by dust

Starting with this type 1 SED, we proceed to model the obscuration of it with dust and gas. In recent times, the type of dust present in type 1 quasars [from the Sloan Digital Sky Survey (SDSS) sample of Richards et al. 2002] has been described as ‘grey’ and alternatively as similar to that of the Small Magellanic Clouds (SMC). Grey dust shows a relatively flat extinction law in the optical and UV, while SMC-type dust is characterized by a steep increase in the extinction at UV wavelengths. Milky Way (MW)-type dust has an UV extinction law which sits in between that of grey and SMC. Richards et al. (2003) and Hopkins et al. (2004) have found the dust in the SDSS type 1 sample to be closest to that found in the SMC, while Czerny et al. (2004) and Gaskell et al. (2004) argued that extinction curves flatten in the UV (grey dust). However, Willott (2005) has shown this implication of grey dust to be a selection effect. The main selection effect arises from measuring dust reddening from a composite constructed from a flux-limited sample: the UV part of the composite is constructed from the higher redshift quasars, which will only make the sample if they are intrinsically brighter and have very little extinction due to dust. This leads to a negative correlation between $E(B - V)$ and redshift, so that different parts of the composite spectrum are dominated by quasars with different amounts of dust. The result is a derived dust extinction curve which appears to flatten in the UV (see Willott 2005, for more details).

We use the extinction curves from Pei (1992), with the extinction at (observed) $24\ \mu\text{m}$ given in terms of the extinction at rest-frame visual band, A_V , and a particular dust type. The quasars in the SDSS sample are all type 1s, and the dust discussed in the above paragraph is not necessarily associated with the torus. The SMC has subsolar metallicity, while the nuclear regions around quasars have solar or supersolar metallicities, with dust properties probably closer to the MW. The effects of different types of dust are secondary compared to the gas-to-dust ratio, yet important, so we assume MW-dust in Fig. 1 and discuss the effect of Large Magellanic Cloud (LMC) and SMC-type dust in Section 2.5 and in Fig. 2. Dust extinction is small at rest frame $24\ \mu\text{m}$, yet the extreme obscuring columns present in the torus and the large redshifts discussed here mean that dust obscuration is not necessarily negligible at observed $24\ \mu\text{m}$. In particular, the $9.7\text{-}\mu\text{m}$ silicate absorption feature can have a very important effect on observed $24\text{-}\mu\text{m}$ flux density at $z \sim 1.5$ (and this feature is significantly deeper for SMC-type dust than for MW-type dust; Fig. 2).

2.3 X-ray absorption

In contrast to dust extinction, which increases with redshift, X-ray absorption decreases at higher frequencies, meaning X-ray samples benefit from a ‘negative K -correction’. The X-ray spectra of heavily absorbed quasars can only be modelled correctly via Monte Carlo simulations, so we use the models of Wilman & Fabian (1999). These assume the intrinsic unabsorbed type 1 spectrum mentioned




Figure 1. Hard X-ray flux ($S_{2-10\text{ keV}}$) versus $24\text{-}\mu\text{m}$ flux density ($S_{24\ \mu\text{m}}$) as a function of redshift for the model quasar described in Section 2. The tracks have the redshifts labelled at the positions marked with black dots at $z = 0.5, 1, 2, 3$ and 5 , and they show the effect on the quasar of MW-type dust with two different gas-to-dust ratios: $N_H = 1.9 \times 10^{25}\text{ m}^{-2} \times A_V$ (MW, top two lines) and $N_H = 5 \times 10^{26}\text{ m}^{-2} \times A_V$ (bottom two lines). Two tracks are drawn for each gas-to-dust ratio, $A_V = 5$ and 50 for the MW gas-to-dust ratio and $A_V = 5$ and 30 for the higher gas-to-dust ratio. The legend shows the combination of A_V and N_H [in $\log_{10}(N_H/\text{m}^{-2})$] for each track. For the larger A_V , the $9.7\text{-}\mu\text{m}$ silicate absorption feature can be seen as a ‘meander’. The dashed straight lines mark the limits for deep surveys covering large areas ($\sim 4\text{--}5\text{ deg}^2$), while the solid straight lines show the limits for the GOODS fields, ultra-deep surveys which cover $\sim 0.1\text{ deg}^2$. As an example of the large area *Spitzer* surveys, we have taken the flux density limit for $24\ \mu\text{m}$ from the FLS (flux limit from Martínez-Sansigre et al. 2005 catalogue from Fadda et al. 2006), while the X-ray flux is taken from the ASCA survey of Ueda et al. (1999). For the deeper GOODS surveys, the $24\text{-}\mu\text{m}$ flux density limit is as quoted by Treister et al. (2006) and the X-ray flux limit is from Alexander et al. (2003). We note that the latter is actually in the $2\text{--}8\text{ keV}$ band (not $2\text{--}10\text{ keV}$ quoted here) but this relatively small difference does not alter our argument. Finally, the dotted lines are for the SXDF limit, a deep survey which has $\sim 1\text{ deg}^2$ of deep X-ray data with $24\text{-}\mu\text{m}$ data from SWIRE (Lonsdale et al. 2003; Ueda et al., in preparation).

earlier, and assume an Fe abundance five times solar, which provides a better fit to the hard X-ray background. The particular models which we use here ignore the reflection component of X-ray spectra, as it has little effect on our discussion, and have a 2 per cent scattered component. The obscuring column of hydrogen, responsible for both types of absorption, is given here as a function of A_V by assuming a gas-to-dust ratio.

The MW is known to have a gas-to-dust ratio such that $N_H = 1.9 \times 10^{25}\text{ m}^{-2} \times A_V$ (Bohlin, Savage & Drake 1978; note this is per magnitude of extinction in V band), while the SMC has $N_H = 1.8 \times 10^{26}\text{ m}^{-2} \times A_V$ (Bouchet et al. 1985). Many AGN have been found to have higher gas-to-dust ratios than the MW (Maiolino et al. 2001; Willott et al. 2004). We therefore choose two representative values for the gas-to-dust ratio. The MW value is sometimes used as a ‘standard’ gas-to-dust ratio. To show the effects of higher gas-to-dust ratios, we note that, for example, NGC 3281 has a value




Figure 2. Curves showing the ratio of transmission for two dust types (LMC and SMC) over the transmission for MW-type dust, using the dust models of Pei (1992), and given two values of A_V . The transmission is here defined as $T = 10^{-A_\lambda/2.5}$. We use $\lambda = 23.7 \mu\text{m}/(1+z)$, the central wavelength of the MIPS instrument at $24 \mu\text{m}$, accounting for the redshift. A_λ is parametrized as a function of A_V and is dependent on the type of dust. We plot and label curves for $A_V = 1$ and 10 , and include LMC as well as SMC-type dust for comparative purposes. The different extinction curves show that, given an A_V , the different types of dust would have significantly different mid-infrared properties, and in particular, type 2 quasars with SMC-type dust will be more difficult to detect in a $24\text{-}\mu\text{m}$ survey than those with MW dust. At $z \gtrsim 3$, the difference becomes small, but around $z \sim 1.5$ SMC-type dust has very deep silicate absorption feature which would make type 2s drop out of flux-limited surveys.

of $N_H = 3.2 \times 10^{26} \text{ m}^{-2} \times A_V$ (Simpson 1998b), while Urrutia et al. (2005) study one object with $N_H \sim 7.7 \times 10^{26} \text{ m}^{-2} \times A_V$ [FTM1004+1229, assuming a factor of $R = 3.0$, the average between the SMC and MW factors, to convert between $E(B - V)$ and A_V]. We therefore choose $N_H = 5 \times 10^{26} \text{ m}^{-2} \times A_V$ as a realistic higher value to illustrate the effect of different gas-to-dust ratios. This is not claiming we encompass the whole range in gas-to-dust ratios, but is chosen to illustrate how mid-infrared and X-ray selection are affected by variations in gas-to-dust ratio. The higher value of gas-to-dust ratio $N_H = 5 \times 10^{26} \text{ m}^{-2} \times A_V$ is high enough for an $A_V = 20$ type 2 quasar to be Compton thick.

2.4 Comparison

Fig. 1 shows the X-ray flux in the $2\text{--}10 \text{ keV}$ band ($S_{2\text{--}10 \text{ keV}}$) versus the $24\text{-}\mu\text{m}$ flux density ($S_{24 \mu\text{m}}$) for type 2 quasars with a range in A_V . MW-type dust is used to obscure the $24\text{-}\mu\text{m}$ emission, and two different gas-to-dust ratios are used. The top set of curves are for a MW gas-to-dust ratio ($N_H = 1.9 \times 10^{25} \text{ m}^{-2} \times A_V$), while the bottom curves are for $N_H = 5 \times 10^{26} \text{ m}^{-2} \times A_V$. For the higher value of A_V , the $9.7\text{-}\mu\text{m}$ silicate absorption feature can be seen as a ‘meander’. The caption in Fig. 1 details the survey limits used for comparative purposes.

The effect of dust at observed $24 \mu\text{m}$ is small in the range $5 \lesssim A_V \lesssim 50$. However, it is important to note that at $A_V \sim 50$, type 2s will start to drop out of $24\text{-}\mu\text{m}$ selection (for the larger area surveys) at $z \sim 1.5$ (due to the silicate absorption) and at $z \gtrsim 2.5$ due to significant dust extinction at the shorter rest-frame wavelengths. For larger values of A_V , the width of the silicate absorption line means type 2s in the range $1.3 \leq z \leq 1.7$ drop out of the sample. As long as the A_V is < 50 , the model type 2 quasar would be detectable by the large area $24\text{-}\mu\text{m}$ surveys at $z \lesssim 3$ [First Look Survey (FLS) and

Spitzer Wide-area Infrared Extragalactic Survey (SWIRE) surveys: Fadda et al. 2006 and Lonsdale et al. 2003, respectively].

In the $\sim 5 \text{ deg}^2$ X-ray ASCA survey (Ueda et al. 1999) with flux limit $10^{-16} \text{ W m}^{-2}$, the model type 2 quasars with $N_H \lesssim 2 \times 10^{27} \text{ m}^{-2}$ are hardly affected by the absorption, and would be detected at $z \lesssim 2$. For $N_H \sim 3 \times 10^{27} \text{ m}^{-2}$, only $z \lesssim 1.5$ type 2s are detected, and the ASCA survey would not have the sensitivity to detect Compton-thick quasars with $N_H \gtrsim 1 \times 10^{28} \text{ m}^{-2}$ above $z = 0.5$.

Since $24\text{-}\mu\text{m}$ selection becomes poor for objects with $A_V \gtrsim 50$ and X-ray selection fails for objects with $N_H \gtrsim 10^{24} \text{ m}^{-2}$, we can see clearly how the gas-to-dust ratio determines which of the two selections is better. For surveys covering a large area ($\sim 4\text{--}5 \text{ deg}^2$), $24\text{-}\mu\text{m}$ selection will be able to pick out type 2s with $N_H \geq 10^{28} \text{ m}^{-2}$ as long as they had $A_V \lesssim 50$. Therefore, for AGN with gas-to-dust ratios like that of the MW or higher, $24\text{-}\mu\text{m}$ selection would fare better than X-ray selection in these surveys. However, we do not know exactly the selection biases in measuring these gas-to-dust ratios, so the possibility of large numbers of AGN with gas-to-dust ratios lower than the MW is not excluded.

For such a gas-to-dust ratio, significantly lower than the MW (e.g. a tenth of the MW value), only quasars with $A_V \gtrsim 50$ would be classified as type 2s by X-ray observations. An extreme $A_V \gtrsim 5000$ would be required for the quasars to be Compton thick. Such objects would not be detected by the larger area mid-infrared samples, and therefore X-ray selection would be more sensitive.

This model type 2 would be detected in the Great Observatories Origins Deep Survey (GOODS) fields at the whole range of redshifts concerned here, both at $24 \mu\text{m}$ and at $2\text{--}10 \text{ keV}$ (survey fluxes from Alexander et al. 2003 and Treister et al. 2006, for X-ray and $24 \mu\text{m}$, respectively). Therefore, although the Compton-thick quasars have not been detected in the ASCA survey, they are, in theory X-ray detectable in the GOODS survey. The reason they have not been found in large numbers is probably an issue of space density, as our model type 2 is rare and large areas are required to find significant numbers of these objects. GOODS therefore has the required sensitivity to find hard X-ray-selected Compton-thick type 2 quasars at all redshifts $z \lesssim 5$, but they do not cover the area required for this.

Since the hard X-ray background is dominated by $z = 0.5\text{--}1$ Seyfert 2s, and since Seyferts have a higher space density than quasars, a large area is not as crucial. However, Seyfert 2s will have typical fluxes $\lesssim 10\text{--}100$ times fainter than our model type 2 quasar, and therefore, at all redshifts Seyfert 2s with $N_H \geq 10^{28} \text{ m}^{-2}$ will have $S_X \lesssim 10^{-19} \text{--} 10^{-18} \text{ W m}^{-2}$. This was, of course, already known, as the deep X-ray surveys in GOODS have not been able to find a population of Seyfert 2s large enough to account for the hard X-ray background. This has been explained by a bias against the most heavily obscured AGN both by X-ray observations and optical spectroscopy (Treister et al. 2004), and our argument is consistent with this.

Thus, to find all type 2 quasars (even the Compton-thick ones), one requires a survey with a $2\text{--}10 \text{ keV}$ flux limit of about $10^{-18} \text{ W m}^{-2}$ over a reasonably large area. Therefore, the Subaru XMM-Newton Deep Field (SXDF) survey (horizontal dotted line; Ueda et al., in preparation), which covers $\sim 1 \text{ deg}^2$ to this depth, should be able to find X-ray-selected Compton-thick quasars. Comparing to the SWIRE depth, we can see that in the region of the SXDF, X-ray selection would be more sensitive than $24\text{-}\mu\text{m}$ selection, while still being able to find powerful type 2 quasars.

For Compton-thick objects, the X-ray selection function is almost flat for $0.5 \lesssim z \lesssim 2$. The redshift distribution of Compton-thick type 2s in a survey deep enough to find them would therefore

be dominated by the volume of the survey as a function of redshift and the evolution of the luminosity function. This means that X-ray surveys deep enough to be sensitive to Compton-thick type 2 quasars should find a wealth of $1 \lesssim z \lesssim 2$ objects. Coincidentally, this would be complementary to 24- μ m surveys, as the mid-infrared surveys would struggle to find heavily obscured type 2s ($A_V \gtrsim 50$), especially in the range around $z \sim 1.5$.

2.5 Dust type

A second-order effect for mid-infrared selection is the type of dust obscuring the AGN, which is particularly important in the range $1.2 \lesssim z \lesssim 1.8$ due to the varying depth of the silicate absorption feature. Fig. 2 shows the ratio of transmission (see figure caption), at observed 24 μ m, for MW- and SMC-type dust. It shows clearly that for $z \lesssim 3$ SMC-type dust will cause more absorption in the mid-infrared, in particular around 9.7- μ m rest frame ($z \sim 1.5$). Thus if the dust obscuring type 2s is more like the dust in the SMC than in the MW, then mid-infrared selection might perform slightly worse than expected from Fig. 1. As mentioned earlier, the dust reddening type 1 quasars has been found to be more similar to SMC dust, but that does not necessarily mean that the dust present in the obscuring torus is also SMC-type dust.

2.6 Orientation of the quasar

The analysis is mainly concerned with 24- μ m selection of objects at $z \lesssim 5$, so we are seeing light reprocessed by dust in the torus at a range of temperatures, $T = 120\text{--}725$ K. We now use the results of a radiative transfer code by Granato & Danese (1994) to investigate the orientation dependence of observed 24- μ m emission. In Fig. 3, we quantify the fraction of the bolometric luminosity, L_{bol} , seen at 24 μ m as a function of redshift. For a pole-on unreddened source (necessarily a type 1), this fraction is approximately constant between $2 \lesssim z \lesssim 5$, while for an edge-on source (a type 2 with $A_V \approx 50$), the fraction is similar to that of a type 1 and approximately constant in the range $2 \lesssim z \lesssim 4$. At higher values of z , the observed 24- μ m corresponds to emitted wavelengths ($\lesssim 5 \mu$ m) at which the dust is becoming progressively more opaque. Around $z \sim 1.5$ the silicate feature is seen in emission by the type 1 and absorption by the type 2, so the observed 24- μ m emission is not a very good indicator of L_{bol} . We can therefore conclude that 24- μ m is a good indicator of bolometric luminosity at $2 \lesssim z \lesssim 4$ and, as shown in Fig. 3, this result is not strongly dependent on the assumed size of the torus.

2.7 Summary

To summarize, we have discussed the advantages of mid-infrared selection over hard X-ray selection. For gas-to-dust ratios similar to those found in the MW, X-ray selection is superior, and can probe further down the luminosity function. However, many AGN have gas-to-dust ratios significantly larger than the MW. We have seen in this section how type 2 quasars with $N_H = 5 \times 10^{26} \text{ m}^{-2} \times A_V$ would naturally drop out of existing large-area X-ray samples without requiring extreme optical obscuration ($A_V \gtrsim 20$ would be enough). We have seen that hard X-ray selected samples are in principle able to find even Compton-thick type 2s, but they require a combination of depth and area that is only just becoming available. We have also seen that dust-type is important, but mainly in the range $1 \lesssim z \lesssim 2$ (due to the 9.7- μ m silicate absorption feature being more pronounced for certain dust types). Using the results of a radiative




Figure 3. Estimate of λF_λ at $\lambda = 24 \mu\text{m}$ (observed) as a fraction of F_{bol} for two Granato & Danese (1994) models with the following parameters: the density has no radial dependence ($\beta = 0$), a covering fraction of 0.8, ratio of outer radius r_m to inner radius of torus r_o (where $r_o \sim 1$ pc), $r_m/r_o = 100$ (top two curves) and $r_m/r_o = 1000$ (bottom two curves). The equatorial optical depth at 0.3 μm , τ_e is 80 (corresponding to $A_V \approx 50$). The dotted line is for an AGN viewed pole-on (a type 1) while the solid line is for an edge-on view (type 2). The reason why the curves only have a few points is that the Granato & Danese (1994) models have few data points at the relevant range of wavelengths, and therefore further smoothing of the curves would be an overinterpretation. This shows that the 24- μ m flux is roughly a constant fraction of F_{bol} between $2 \lesssim z \lesssim 4$ for both type 1 and type 2s. In this range of redshifts, 24 μ m is therefore a good wavelength to select AGN independent of orientation. Once again we see the problems caused by the silicate absorption at $z \sim 1.5$. At very high redshifts ($z \gtrsim 4$), observed 24 μ m is now a relatively short wavelength in the rest frame, meaning obscuration by dust starts to have an increasingly large effect. The curves for both r_m/r_o ratios are very similar, except for a normalization, showing this argument holds for a range of torus sizes.

transfer code, we have also seen that (observed) 24- μ m luminosity is a constant tracer of type 2 bolometric luminosity in the range $2 \lesssim z \lesssim 4$. Therefore, 24- μ m selection is a natural starting place to look for the population of high-redshift type 2 quasars missing from X-ray surveys.

3 SELECTION CRITERIA

Our aim was to find the elusive high-redshift type 2 population around the peak in the quasar activity, $z \gtrsim 2$ (e.g. Wolf et al. 2003), and for this we used the following selection criteria:

- (i) $S_{24 \mu\text{m}} > 300 \mu\text{Jy}$
- (ii) $S_{3.6 \mu\text{m}} \leq 45 \mu\text{Jy}$
- (iii) $350 \mu\text{Jy} \leq S_{1.4 \text{ GHz}} \leq 2 \text{ mJy}$.

3.1 Data sets

The work described in this paper was all done in the *Spitzer* extragalactic FLS. Three separate catalogues were cross-matched for the initial selection: the IRAC 3.6 μ m (Lacy et al. 2005b) and the MIPS 24 μ m (Marleau et al. 2004; Fadda et al. 2006) from the *Spitzer* FLS and the 1.4-GHz catalogue from the NRAO VLA (Condon et al. 2003). The flux limits used for 3.6 μ m, 24 μ m and 1.4 GHz were 20, 300 and 100 μJy , respectively.

Over the area with coverage in all three bands (3.8 deg^2), the radio and 24- μ m catalogues were matched, to select all sources detected at 24 μ m and within the radio flux cuts. The chosen sources were then contrasted with the 3.6- μ m catalogue, to obtain their flux densities

in this IRAC channel. However, objects were not required to be detected at 3.6 μm to be kept in the candidate list. All matching was done with a 2.5-arcsec tolerance to allow for positional offsets between catalogues.

In addition, in this paper we use tabulate R -band magnitudes from data from the 4-m Mayall Telescope Kitt Peak National Observatory (Fadda et al. 2004), and we also make use of a Westerbork Synthesis Radio Telescope (WSRT) deep survey at 1.4 GHz in the *Spitzer* FLS (Morganti et al. 2004). However, these two data sets were not used to select the candidate type 2 quasars.

3.2 Mid-infrared criteria

The mid-infrared criterion $S_{24\mu\text{m}} > 300 \mu\text{Jy}$ was chosen to select a flux-density-limited sample of active galaxies. At lower redshifts, a 24- μm selected sample would be dominated by starbursting galaxies, but combined with the 3.6- μm selection (which ensured high-redshift objects, see below), this 24- μm flux limit should yield powerful AGN. The $S_{24\mu\text{m}} > 300 \mu\text{Jy}$ criterion is the flux-density limit from a preliminary catalogue, and is actually a 7σ limit in the 24- μm final catalogue from the *Spitzer* FLS data (Marleau et al. 2004; Fadda et al. 2006). At $z = 2$, this corresponds to an emitted 8- μm luminosity $L_{8\mu\text{m}} = 10^{24.4} \text{ W Hz}^{-1}$. Assuming a typical type 1 SED (Rowan-Robinson 1995), this is $L_B = 10^{23.2} \text{ W Hz}^{-1}$ or $M_B = -23.8$. At $z = 2$, the break in the quasar luminosity function, L_{quasar}^* , has a value of $M_B = -25.7$ (Croom et al. 2004, ; with pure luminosity evolution), so our 24- μm selection will select quasars $\gtrsim 0.2 L_{\text{quasar}}^*$ at $z = 2$, and more luminous quasars at higher redshifts. As explained in Section 2, the selection becomes more complicated at $z < 2$ due to the silicate absorption, so we restrict our discussion to $z \geq 2$. The peak of the quasar activity occurred around $z \geq 2$ and by targeting the quasars around the ‘break’ we are sensitive to the part of the population that contributes most of the luminosity density.

Quasars are considered as being type 2 if $A_V \gtrsim 5$ (Simpson et al. 1999), and such an amount of extinction will make the observed near-infrared emission of type 2s much fainter than that of type 1s. The 3.6- μm criterion was therefore chosen to remove naked (type 1) quasars as well as lower redshift ($z \lesssim 1.4$) type 2s. At $z \geq 2$, the detected 3.6- μm flux density corresponds to light emitted at $\lambda \leq 1.2 \mu\text{m}$ so dust extinction ensures that type 2 quasars are much fainter than type 1 quasars, even for a moderate A_V . Indeed, the $S_{3.6\mu\text{m}}$ emission is likely to be dominated by starlight for $A_V \gtrsim 10$, and since light at 3.6- μm is dominated by the old stellar population, there will be an $S_{3.6\mu\text{m}} - z$ correlation, analogous to the $K - z$ relation for radio galaxies (Jarvis et al. 2001b; Willott et al. 2003b). A typical host galaxy for a $z = 2$ radio-quiet quasar is the progenitor of a $\sim 2L_{\text{gal}}^*$ (Kukula et al. 2001) galaxy in the local Universe, so we adapt the $K - z$ relation (which corresponds to $\sim 3L_{\text{gal}}^*$ hosts for radio galaxies) to $\sim 2L_{\text{gal}}^*$ hosts at 3.6 μm . To do this, we use the models of Bruzual & Charlot (2003) to form an elliptical galaxy at $z = 10$, assign it a (stellar) mass and allow it to evolve passively. This allows us to predict the 3.6- μm flux as a function of redshift, for an elliptical galaxy of this given mass, and hence obtain a rough photometric redshift. The stellar mass of the galaxy was set to $3.8 \times 10^{11} M_\odot$, and in Martínez-Sansigre et al. (2005) we quoted this as corresponding to $\sim 2L_{\text{gal}}^*$. If we assume a Salpeter IMF (initial mass function) (Salpeter 1955) and take $M_K = -24.3$ from the fitting of Schechter functions by (Cole et al. 2001), then our assumed stellar mass translates to a $2.6L_{\text{gal}}^*$ galaxy. This is the host (elliptical) galaxy luminosity assumed to obtain the rough photometric redshifts in Martínez-Sansigre et al. (2005) and in this paper. The




Figure 4. Mid-infrared selection criteria plotted on observed SEDs. The 24- μm criterion is shown as a lower limit, and the 3.6- μm criterion as an upper limit. The solid lines represent quasars at $z = 2$ bright enough to make the 24- μm flux cut. The upper solid line is a type 1 quasar (SED from Rowan-Robinson 1995), while the lower solid line is a type 2 with $A_V = 5$. A type 1 quasar just bright enough to make the 24- μm cut will be too bright at 3.6 μm (once the host galaxy contribution is added). The dashed SED represents a $2.6L^*$ elliptical galaxy at $z = 1.4$ (from Bruzual & Charlot 2003), where it becomes just faint enough to make it fall below the 3.6- μm cut. The dotted line is the ULIRG NGC 7714 redshifted to $z = 0.4$ (SED from Brandl et al. 2004), which would meet our mid-infrared criteria if it lay between $z = 0.25$ and $z = 0.4$.

criterion $S_{3.6\mu\text{m}} \leq 45 \mu\text{Jy}$ corresponds to a limiting photometric redshift $z_{\text{phot}} \gtrsim 1.4$. This was chosen to target $z \sim 2$ type 2 quasars, while allowing for scatter in the crude photometric redshift estimation and filtering out type 1 quasars and low-redshift contaminants like radio galaxies.

The combination of these criteria is illustrated in Fig. 4. It shows the observed SEDs of two $z = 2$ quasars (model from Rowan-Robinson 1995), a type 1 and a type 2 (the latter obscured by $A_V = 5$, dust model from Pei 1992). A $2.6L^*$ elliptical galaxy from Bruzual & Charlot (2003) is also plotted at $z = 1.4$, showing how the 3.6- μm criterion serves also as a crude photometric redshift cut.

3.3 Radio criterion

The radio selection criteria are added to ensure that the candidates are quasars rather than starburst galaxies. The 3.6–24- μm ‘colour’ demanded by our criteria can be achieved by lower redshift ($z < 1$) ultraluminous infrared galaxies (ULIRGs). For example, NGC 7714 would satisfy these criteria if it were at $z = 0.25$ – 0.4 (see fig. 4, SED from Brandl et al. 2004) and the more luminous ULIRG Arp 220 if it were at $z = 0.5$ – 0.7 . In the absence of AGN radio activity, the radio flux densities of such objects would follow the far-infrared/radio correlation (Condon 1992), and hence would have values of $S_{1.4\text{GHz}} \sim 50$ (for ULIRGs at $z \sim 0.3$) and $500 \mu\text{Jy}$ (for more powerful ULIRGs at $z \sim 0.6$).

To avoid such starburst contaminants, we chose a lower limit on $S_{1.4\text{GHz}}$ which also excludes higher redshift (submm selected) starburst galaxies without the benefit of strong gravitational lensing (e.g. Chapman et al. 2005). For a $z \sim 0.6$ ULIRG to have $S_{1.4\text{GHz}} \geq 350 \mu\text{Jy}$, it would require a far-infrared luminosity of $L_{\text{FIR}} \gtrsim 1 \times 10^{12} L_\odot$. Objects with this far-infrared luminosity are rare in space density (Sanders & Mirabel 1996), and the comoving volume probed at $z \sim 0.6$ by 3.8 deg^2 is such that $\lesssim 1$ contaminants would be expected in our sample.

Table 1. Basic data on the 21 type 2 quasars in our sample. The J2000.0 positions are from the *Spitzer* FLS radio catalogue (Condon et al. 2003). The MIPS 24- μm flux density $S_{24\mu\text{m}}$ is obtained by point spread function (PSF) fitting (Marleau et al. 2004) as all objects (except AMS16) are point sources at the ~ 6 arcsec resolution of the MIPS observations, with positional errors ~ 1 arcsec. It has a typical error of ± 10 –15 per cent. The 1.4-GHz flux density is the integrated value (in μJy) from the radio catalogue. The IRAC 3.6- and 4.5- μm flux densities are measured in 5-arcsec diameter apertures (Lacy et al. 2005b) and have typical errors of ± 10 per cent. The *R*-band magnitudes (Vega) are measured using a 3-arcsec diameter aperture from the images of the 4-m Mayall Telescope Kitt Peak National Observatory (Fadda et al. 2004), with typical errors of 0.03 for objects with magnitude 22–23, 0.10 for objects with 23–24 and 0.25 mag for those with magnitudes of 25.

Name	RA (J2000)	Dec.	$S_{24\mu\text{m}}$ (μJy)	$S_{1.4\text{GHz}}$ (μJy)	$S_{3.6\mu\text{m}}$ (μJy)	$S_{4.5\mu\text{m}}$ (μJy)	m_R	z_{phot}	z_{spec}
AMS01	17 13 11.17	+59 55 51.5	536	490	25.0	30.0	25.1	2.1	
AMS02	17 13 15.88	+60 02 34.2	294	1184	44.5	61.0	25.3	1.4	
AMS03	17 13 40.19	+59 27 45.8	500	1986	16.0	28.0	24.4	3.1	2.698 ^a
AMS04	17 13 40.62	+59 49 17.1	828	536	18.0	22.0	22.96	2.8	1.782
AMS05	17 13 42.77	+59 39 20.2	1769	1038	34.7	61.4	25.2	1.7	2.017
AMS06	17 13 43.91	+59 57 14.6	969	444	<20	<25	≥ 25.3	≥ 2.5	1.76 ^b
AMS07	17 14 02.25	+59 48 28.8	503	354	37.9	47.8	24.7	1.5	
AMS08	17 14 29.67	+59 32 33.5	792	655	41.6	46.3	22.48	1.4	1.979
AMS09	17 14 34.87	+58 56 46.4	685	426	25.2	<25	≥ 25.3	2.1	
AMS10	17 16 20.08	+59 40 26.5	338	1645	39.2	44.7	24.1	1.5	
AMS11	17 18 21.33	+59 40 27.1	442	356	32.4	55.6	24.7	1.7	
AMS12	17 18 22.65	+59 01 54.3	518	946	25.24	<25	23.77	2.0	2.767
AMS13	17 18 44.40	+59 20 00.8	4196	1888	24.7	49.1	22.89	2.1	1.974 ^c
AMS14	17 18 45.47	+58 51 22.5	937	469	8.0	15.0	23.67	4.6	1.794
AMS15	17 18 56.93	+59 03 25.0	371	440	16.0	16.0	≥ 25.3	3.0	Not observed ^d
AMS16	17 19 42.07	+58 47 08.9	788	390	18.0	21.0	23.07	2.8	4.169
AMS17	17 20 45.17	+58 52 21.3	1134	615	10.0	15.0	23.80	3.9	3.137 ^a
AMS18	17 20 46.32	+60 02 29.6	925	390	<20	<25	23.53	≥ 2.5	1.017
AMS19	17 20 48.00	+59 43 20.7	1433	822	<20	26.9	≥ 25.3	≥ 2.5	2.25 ^b
AMS20	17 20 59.10	+59 17 50.5	492	1268	45.2	64.9	23.80	1.4	^e
AMS21	17 21 20.09	+59 03 48.6	720	449	25.2	38.6	24.2	2.1	

^aThese objects have redshifts identified from a single line, which we identified as Ly α : in AMS03, from the amount of structure in the line, and in AMS17 from the fainter continuum blueward of the line.

^bThese two objects have redshifts from *Spitzer*–IRS spectroscopy from Weedman et al. (2006), who give two values for the redshifts. Here, we show the mean of the two values.

^cThis object was observed with *Spitzer*–IRS by Yan et al. (2005) who measured a redshift of $z = 2.1 \pm 0.1$ in good agreement with our value.

^dAs explained in Table 2, AMS15 has not really been observed spectroscopically.

^eAMS20 shows faint red continuum but no emission lines. Note there were errors in the redshifts of AMS14 and AMS18 presented in table 1 of Martínez-Sansigre et al. (2005) and these have been corrected here, together with minor corrections to several other redshifts.

An upper limit on $S_{1.4\text{GHz}}$ was also used to filter out the radio-loud objects, whose extended jets might complicate interpretation. The radio flux-density cut introduced here will select the radio-bright end of the radio-quiet quasar population (Cirasuolo et al. 2003), so these objects can be described as radio intermediate. In addition, the $S_{3.6\mu\text{m}}$ criterion includes objects without IRAC detections which we expect to have the highest redshifts. Since the 24- μm positions are accurate to ~ 1 arcsec, the FLS radio positions (Condon et al. 2003), accurate to ~ 0.5 arcsec, were also better for spectroscopic follow-up.

The two simple infrared criteria together with radio selection were chosen over an IRAC colour requirement because the latter criterion would have selected type 2s with a narrow range in A_V and z . Our three criteria represent a simple way of selecting type 2 quasars and allow the modelling necessary to estimate the quasar fraction at high redshift (Martínez-Sansigre et al. 2005).

After these selection criteria, we were left with 21 candidate high-redshift type 2 quasars (see Table 1 and Fig. 5), which were followed up spectroscopically (as described in Section 4). Note that all our candidates were originally selected from the preliminary 3.6- and 24- μm catalogues, but have had their fluxes slightly revised and the flux densities quoted are the final ones. This explains why two candidates are slightly outside the selection boundaries. We also note that although the *R*-band (Vega) magnitudes are not part of our

selection criteria, they are consistent with optically obscured quasars in high-redshift host galaxies. At $z = 1.4$, our model elliptical galaxy would have $m_R = 23.3$.

4 OBSERVATIONS AND DATA REDUCTION

The 21 candidates were observed with both arms of the ISIS instrument at the William Herschel Telescope in 2004 July and 2005 April (see Table 2). Low-resolution long-slit spectroscopy was performed across the entire visible band (3200–9300 Å) using the gratings R158B and R158R, and the EEV12 and Marconi detectors. The 6100-Å dichroic was used in 2004 July, while in 2005 April we used the new 5300-Å dichroic. Using slit widths of 1.5–2.0 arcsec, we obtained spectral resolutions of 12–16 Å in both arms. Bias, flat, arc, sky-flat and standard-star frames were all taken at the beginning and end of the night to ensure reasonable calibration.

The slit was placed on the radio positions, without optical identification, by ‘blindly’ offsetting from a nearby guide star (e.g. Rawlings, Eales & Warren 1990). Whenever another radio source was in the field, the slit angle was chosen to go through the source as well as our candidate. When no other radio sources were available, bright sources from the IRAC 3.6- μm catalogue were placed in the slit. This ensured multiple objects were detected in the slit, improving the astrometry and identification of our sources in the 2D

Table 2. Summary of the spectroscopic observations. Most of the objects were observed in 2004 July, and the spectroscopy was completed in 2005 April. The objects were all observed with airmass < 1.5 (except AMS05), with good seeing (in the range 0.7–1.0 arcsec). As described in Section 4, the PAs of the slit were generally chosen to go through any other radio source in the field, or any bright 3.6- μm source, to help the astrometry and identification. In some cases neither was available, nor the airmass was high, so the PA was chosen to match the parallactic angle and therefore minimize dispersion of the blue light.

Name	RA (J2000)	Dec.	Date observed	Airmass	PA ($^{\circ}$)	Slit (arcsec)
AMS01	17 13 11.17	+59 55 51.5	04/07/15	1.17	103	1.5
AMS02	17 13 15.88	+60 02 34.2	04/07/14	1.37	149	2.0
AMS03	17 13 40.19	+59 27 45.8	04/07/14	1.17	114	1.5
AMS04	17 13 40.62	+59 49 17.1	05/04/14	1.18	36	2.0
AMS05	17 13 42.77	+59 39 20.2	04/07/14	1.47/1.56 ^a	98	2.0
AMS06	17 13 43.91	+59 57 14.6	04/07/13	1.19	120	2.0
AMS07	17 14 02.25	+59 48 28.8	04/07/16	1.22	19	1.5
AMS08	17 14 29.67	+59 32 33.5	04/07/15	1.32	128	1.5
AMS09	17 14 34.87	+58 56 46.4	04/07/15	1.17	111	1.5
AMS10	17 16 20.08	+59 40 26.5	04/07/14	1.20	114	1.5
AMS11	17 18 21.33	+59 40 27.1	04/07/16	1.19	76	1.0
AMS12	17 18 22.65	+59 01 54.3	04/07/15	1.16	150	1.5
AMS13	17 18 44.40	+59 20 00.8	05/04/14	1.16	186	2.0
AMS14	17 18 45.47	+58 51 22.5	04/07/13	1.20	105	2.0
AMS15	17 18 56.93	+59 03 25.0	05/04/15 ^b	–	–	–
AMS16	17 19 42.07	+58 47 08.9	04/07/16	1.25	141	2.0
AMS17	17 20 45.17	+58 52 21.3	05/04/15	1.27	117	1.5
AMS18	17 20 46.32	+60 02 29.6	04/07/13	1.40	133	2.0
AMS19	17 20 48.00	+59 43 20.7	05/04/14–15	1.17/1.17	170/185	2.0
AMS20	17 20 59.10	+59 17 50.5	04/07/14	1.22	160	2.0
AMS21	17 21 20.09	+59 03 48.6	05/04/15	1.20	84	1.5

^aAMS05 was observed twice on the same night, at different airmasses.

^bAMS15 was targeted in the 2005 April, but the position used was slightly wrong (~ 0.8 arcsec) and therefore we do not consider it to have been observed.

spectra. At high-airmasses (~ 1.4 – 1.5), we observed the objects with slit position angles (PAs) close to the parallactic angle, to minimize atmospheric dispersion of the blue light. Each object was observed for 30 min with a 1–2 arcsec slit width, with one 1800-s exposure in the blue arm and two 900-s exposures in the red arm.

The spectra were all reduced using the IRAF package TWODSPEC LONGSLIT. The wavelength calibration was performed using CuAr+CuNe lamps and flux calibration was obtained from appropriate spectrophotometric stars. The two red frames for each object were reduced separately and then combined to remove cosmic rays. The 1D spectra were extracted separately for the blue and the combined red frames, using the task APSUM. The resulting one- and two-dimensional spectra are plotted in Fig. 6.

AMS01. The 3.6- μm image of this object shows it to be visible, although possibly confused with a galaxy just to the north (N) of it (see Fig. 5). However, in the *R*-band image we see our target is distinguishable as an extremely faint galaxy. The two images clearly show the very red colour of AMS01. There was another radio source of similar flux east (E) of AMS01 and so a PA of 103° was chosen to go through both sources. The spectrum of AMS01 was completely blank in a 30-min exposure.

AMS02. In this case, our target is clearly visible in the IRAC image, but is only just detected in the *R* band. The PA was chosen to go through another radio source, 45-arcsec south-east (SE) of AMS02. The spectrum of our target was once again completely blank in a 30-min exposure.

AMS03. Here we see another possible source of confusion in the 3.6- μm image, but the *R*-band image shows our source to be

very faint but distinct from the brighter source south-west (SW) of it. There is a nearby radio source which is confused in the WSRT images (see Section 8). Here, the PA was chosen to go through this other radio source 18 arcsec E–SE of AMS03. The two-dimensional spectrum shows a faint, double, spatially extended line (~ 1.6 arcsec once deconvolved from the seeing), which we identify as Ly α (the redward line is not N v). The spectrum shows quite a lot of structure (see Fig. 6): there are two Ly α lines, one at 4481 Å and the other at 4512 Å. The redward line has a full width at half-maximum (FWHM) of $\sim 500 \text{ km s}^{-1}$, while the blueward line has a hint of absorption in the spatial centre, and has a greater FWHM ($\sim 1400 \text{ km s}^{-1}$) but still narrow enough to qualify as a type 2 quasar. A possible interpretation is that the type 2 quasar resides in a Ly α halo, and there is a large difference in systemic velocity for the quasar and the halo (or the halo is collapsing, leading to an additional redshift). Another possibility is associated absorption – this is present in small radio sources (van Ojik et al. 1997).

AMS04. This object is very faint at 3.6 μm (18 μJy) but reasonably bright in *R* band ($m_R = 23$). The radio emission around AMS04 is extended towards the north-east (NE). There is another distant radio source, identified at both 3.6 μm and *R* band further to the east. This is possibly associated with another extremely faint galaxy about 10-arcsec away, although the radio and optical positions of the adjacent radio source and the faint galaxy do not match perfectly. We placed the slit to go through both AMS04 and the other radio source. AMS04 shows strong continuum, a spatially extended (and bright) Ly α as well as [C III], [N II] and [C II] lines but nothing was seen associated with the extended radio emission. The Ly α line




Figure 5. Radio contours overlaid over the 3.6- μ m and R -band images for the sample (images from Condon et al. 2003; Fadda et al. 2004; Lacy et al. 2005b). In each pair of overlays, the 3.6- μ m image is on the left-hand side (with the name of the source) and the R band on the right-hand side. Radio contours are plotted at -2σ , 2σ and subsequently at $2^{1/2}\sigma$ up to $2^6\sigma$, where σ is the rms noise of the radio image ($\approx 23 \mu\text{Jy}$). The grey-scale has been chosen to overcome the high dynamical range and highlight faint objects. Faint grey pixels have values three times that of the noise.

has an angular size of ~ 5.5 arcsec. This object has a redshift of 1.782, with all the redshifts from individual lines agreeing well (see Table 3). The fact that this object is not that faint in R band is due to the continuum and narrow lines at the red end of the spectrum.

AMS05. Here, we see again a source which is relatively bright at 3.6 μ m (34 μJy) compared to R band ($m_R = 25.2$). The IRAC image is quite crowded and there is the danger of confusing sources, but we can see in the R band that the radio position falls on a faint galaxy. This object showed an interesting spectrum: C IV is actually as strong as Ly α , and He II is also detected, but there is no continuum in the blue or red arm. All lines are unresolved, and the weakness of the Ly α may be due to it falling in the blue end of the spectrum while having been observed at a high airmass (see Table 2) and away from the parallactic angle.

AMS06. This target is just about visible in the 3.6- μ m images, but it is not bright enough to make it to the IRAC catalogue. It is

undetected in the R band. A PA of 120° was chosen to go through the ‘semimajor axis’ of the marginally resolved radio source. The spectrum was blank at the location of the target source. This could be due to the entire galaxy being very dusty and therefore faint in the optical and even in the infrared (see the Section 9) or it could be a very high redshift source. Recently, a redshift of 1.76 has been obtained from silicate absorption and polycyclic aromatic hydrocarbon (PAH) emission with *Spitzer* IRS (Weedman et al. 2006), confirming this is a very dusty system at high redshift.

AMS07. AMS07 is quite a red object (within our sample) in that the 3.6 μ m is relatively bright, whereas the R band is very faint. There is another radio source 56-arcsec SW of the target, so the slit (with PA 19°) was made to go through both radio sources. The spectrum was blank at the target position.

AMS08. Here, we have a relatively bright source for our sample, both in the IRAC and R bands. A faint (slightly extended) radio




Figure 5 – continued

source, 18-arcsec SE of our source, was visible in the radio images. The slit was made to go through both this source and our target. AMS08 appears in the spectrum, showing Ly α (~ 3.7 -arcsec wide after deconvolving from the seeing), C IV and He II and blue continuum.

AMS09. This is a faint source at $3.6\ \mu\text{m}$ and is undetected in R band. The $3.6\text{-}\mu\text{m}$ image shows only one infrared ID for our radio source, but the R -band image shows there might be a serious confusion issue, as there are several faint sources near the radio position. There were two other radio sources in the field, one close to the N (visible in Fig. 5) and one to the E. We chose to place the slit through the eastern radio source (41-arcsec away) and the PA angle (119°) is actually appropriate to try and catch several of the possible optical IDs. There is barely a hint of continuum from AMS09 in the red two-dimensional spectrum and no hint of the other source.

AMS10. AMS10 is a relatively bright source in the $IRAC$ bands, and has $m_R = 24.1$. The $3.6\text{-}\mu\text{m}$ source is actually confusing two R -band sources, but the radio position is clearly associated with only one of the R -band galaxies. There were no other radio sources in

the field, so we chose the PA to match the parallactic angle. The 2D spectrum of AMS10 is blank.

AMS11. This is another quite red source, relatively bright at $3.6\ \mu\text{m}$ and faint in R band. There are no other radio sources near but the radio emission seems to be slightly extended in the E direction, so we chose the PA so the slit would go through it. AMS11 was not detected by ISIS.

AMS12. Here, we have a faint source in all bands. The PA was chosen to go through a radio source 50-arcsec SE of our target. The spectrum of AMS12 shows several emission lines: Ly α (spatially unresolved), C IV and He II (see Table 3) and faint continuum.

AMS13. AMS13 is a point-like radio source, and is quite faint at $3.6\ \mu\text{m}$, so we matched the PA to the parallactic to minimize dispersion of the blue light (even though the airmass, 1.16, was not particularly high). In the 2D spectrum, however, it is very bright, showing spatially extended Ly α and C IV lines (~ 4.3 - and 3.5 -arcsec wide, respectively) as well as continuum. The continuum explains why it is not that faint in the R band. The optical spectra show AMS13 is at a redshift of 1.974, and Yan et al. (2005) find a redshift




Figure 5 – continued

of 2.1 ± 0.1 from the silicate absorption feature in their *Spitzer*–*IRS* spectroscopy, in agreement with our redshift.

AMS14. This source is extremely faint in the *IRAC* bands (8 μ Jy at 3.6 μ m) making us expect this to be a very high redshift source. A PA of 105° was chosen to go through another faint radio source 10-arcsec E of our target. The spectrum, however, shows a very strong (and spatially resolved) $\text{Ly}\alpha$ line (~ 1.6 -arcsec wide) as well as C IV at a redshift of $z = 1.794$. A mistake in an early wavelength calibration led us to misidentify the lines and the redshift reported in Martínez-Sansigre et al. (2005) is incorrect.

AMS15. This is another very faint source at 3.6 μ m and undetected in *R* band. This is a good candidate for a high-redshift type 2 quasar. However, we used a slightly wrong position (the one published in Martínez-Sansigre et al. 2005 is off by ~ 0.8 arcsec) for follow-up spectroscopy. For this reason we do not consider this object to have been observed.

AMS16. AMS16 is very faint in the *IRAC* bands, although it is not extremely faint in *R* band ($m_R = 23.07$). It has an extended radio structure, whose central position matches the optical and infrared positions. However, the extended emission matches very well two galaxies next to the source, and in the 24- μ m image the three sources are confused, and have a similar structure to the radio emission. There is the possibility that the two foreground galaxies might be lensing AMS16 (see Fig. 5). Two factors therefore influenced the choice of PA in this case. First, the radio emission around AMS16 is extended in the SE to NW direction (diagonally, possibly due to the two galaxies). In addition, there is a radio loud ($S_{1.4\text{ GHz}} = 21.6$ mJy) type 1 quasar about an arcmin NW. By good fortune, the

same angle is required to get the slit through the ‘semimajor axis’ of the extended radio emission, the brighter foreground galaxy and the type 1 quasar. We found $\text{Ly}\alpha$ (~ 2.7 arcsec), N V (~ 1 arcsec) and C IV (spatially unresolved) in AMS16, indicating a redshift of $z = 4.169$. The emission lines of AMS16 are all strong, which explains why it is not that faint in the *R* band. To the NW of AMS16, 58-arcsec away, we found a radio-loud type 1 quasar, at redshift $z = 3.785$ (see Fig. 7).

AMS17. This is another faint source in the *IRAC* bands, suggesting a high redshift, although the *R* band is not particularly faint for this sample, and the images show the optical/infrared positions match perfectly the centre of the radio emission, with no possible sources of confusion nearby. The slit PA was chosen to go through another object, about 59-arcsec SE of our target. AMS17 is indeed at high redshift, as it shows one narrow line in the blue spectrum (which we identify as $\text{Ly}\alpha$ at $z = 3.137$) and faint continuum redwards of the line. The $\text{Ly}\alpha$ line is extended ~ 1.6 -arcsec wide in the spatial direction, once the seeing is deconvolved.

AMS18. Here is a source which is undetected in the *IRAC* catalogues, which suggests a high redshift. The PA was chosen to go through a bright radio source 57-arcsec SE. The 2D spectrum shows AMS18 has several faint emission lines, which we identify as [C II], Mg II and [O II] at $z = 1.017$, so it is at a lower redshift than first expected from the crude photometric redshift. Note the weakness of the [O II] line, which suggests a faint AGN rather than something similar to a radio galaxy. The same wavelength calibration error discussed for AMS14 led to the redshift of AMS18 being incorrect in Martínez-Sansigre et al. (2005). This is probably a less-luminous




Figure 6. Optical spectra of objects in the sample which showed emission lines. The emission lines from which the redshifts were obtained are labelled, while tentative identifications of lines which have not used for the redshifts are labelled with a question mark. The corresponding two-dimensional spectra are shown underneath the one-dimensional ones as a visual help to find the emission lines. They have approximately the same scale in the horizontal direction as the one-dimensional spectra. The grey-scale has been chosen to bring out low signal-to-noise ratio lines and can lead to visual artefacts such as the strange change in continuum strength of AMS08. For the two objects observed in 2005 April (AMS04 and AMS13), the pixel scale in the dispersion axis was not the same for the blue and red arms, and so the two-dimensional spectra are not combined, just overlaid on each other. Particularly, strong residual sky lines have been labelled in the one-dimensional spectra and are clearly visible in the two-dimensional ones. In AMS04, the sky lines are close to the [N II] and [C II] lines, making these two lines hard to detect in the grey-scale image. The reader is assured that these two lines are detected and are not due to the residual sky lines.

quasar possibly hosted by a small galaxy, which is how it made it through our selection criteria. It is, however, difficult to judge whether the lines are spatially extended, due to the low signal-to-noise ratio.

AMS19. The PA was chosen to be identical to the parallactic angle, as it is very faint in the near-infrared and the radio emission of AMS19 is point like. This could mean a high redshift, so we tried

to maximize the blue light received. Indeed, in the two-dimensional AMS19 does not even show faint continuum. However, it was also observed by Weedman et al. (2006) with *Spitzer* IRS and is found to be at $z = 2.25$ from its PAH and Silicate features.

AMS20. This is a relatively bright source for our sample, and so is not expected to lie at a particularly high redshift. The slit was placed at an angle so it would go through two fainter radio




Figure 6 – continued

sources NW of AMS20 (19- and 55-arcsec away). The spectrum of AMS20 shows faint red continuum but no lines to allow us to obtain a redshift. Further study is required to determine whether this is a very bright high-redshift type 2, or more similar to AMS18, a faint, lower redshift object. The lack of lines suggests $z \sim 1.3$ – 1.6 .

AMS21. The last source in our sample is relatively faint in at $3.6 \mu\text{m}$ and R band. In this case, the PA was chosen to go through a fainter radio source about 25-arcsec E of the target. However, neither AMS21 nor the other radio source has been detected in the optical spectrum.

5 TYPE 2 SPECTRA

Approximately half of the objects (10 out of 21) showed narrow emission lines and sometimes continuum (see Fig. 6). $\text{Ly}\alpha$ with $\text{FWHM} \lesssim 2000 \text{ km s}^{-1}$ was found in nine objects (with a redshift range $1.78 \leq z \leq 4.17$) and high excitation lines such as C IV (1549 Å) or He II (1640 Å) were also found in several objects (see Table 3). In two cases (AMS03 and AMS17), we only found one emission line, which was taken to be $\text{Ly}\alpha$ by virtue of its extreme equivalent width, blue-absorbed line profile and lack of other lines supportive of alternative identifications. The $\text{Ly}\alpha$ line of AMS03 is particularly spatially extended. One object, AMS18, proved to be at $z = 1.02$, lower than expected from our selection criteria. Only one object showed (very faint) continuum and no lines: AMS20. As mentioned in Section 4, it will require more study to determine whether this object is similar to AMS18, is at $1.3 \lesssim z \lesssim 1.7$, or is at a higher redshift object. If this object were a contaminant, like a $z \sim 0.5$ ULIRG, it would have shown stronger continuum and the $[\text{O II}]$

3727-Å line. It is important to insist on how faint our sources are: all objects, even those with emission lines in the R band or showing some continuum, have $m_R \geq 22.5$ (Vega), and most have $m_R \geq 23$. The rest of the objects show blank spectra, confirming that there are no low-redshift contaminants in our sample. A $2.6L^*$ elliptical at $z = 1.4$ would have a continuum flux density of $1 \times 10^{-21} \text{ W m}^{-2} \text{ Å}^{-1}$, at the limit of our sensitivity. AMS20, with faint red continuum, is therefore consistent with having an elliptical host galaxy, and being somewhere in the spectroscopic desert.

The nature of the blank-spectrum objects is discussed further in Section 9, but we see that their blank optical spectra are consistent with host ellipticals at $z \gtrsim 1.4$. If $24\text{-}\mu\text{m}$ flux density is a good tracer of narrow emission-line strength, then we expect all our sources to have detectable $\text{Ly}\alpha$ lines if they lie at $z \geq 1.7$. In addition, the C IV (1549 Å) line is detected in six out of the nine objects with $\text{Ly}\alpha$, and in the composite spectrum (see Section 7) it has 36 per cent of the flux of $\text{Ly}\alpha$. Hence, some objects in the spectroscopic desert should also have detectable C IV lines. The two blank objects brightest at $24\text{-}\mu\text{m}$ have redshifts from *Spitzer*–*IRS* in the same range as the rest of the optical spectra, where we should have seen $\text{Ly}\alpha$. This suggests that at least some of the blank objects are at redshifts high enough to have narrow emission lines in the optical band, and that their spectra are blank for another reason. This reason could be that they have weaker emission lines, but they would have to be significantly weaker than those of the narrow-line objects, despite having similar $24\text{-}\mu\text{m}$ fluxes. Another alternative is that the narrow emission-line region is itself obscured, as has proven to be the case from the *Spitzer* spectroscopy of AMS06 and AMS19.

Table 3. Table of emission-line properties of the 10 objects with spectroscopic redshifts from our optical spectroscopy. The lines were fitted with Gaussians to find λ_{obs} , FWHM, and the flux. For objects with several lines, the error quoted is the square root of the average variance. For objects with only one line, the error is assumed to be dominated by the fitting of a Gaussian to the line. This in turn is dominated by the spectral resolution and the seeing. In two cases (AMS14 and AMS18), an error in wavelength calibration led to incorrect identification of lines and redshifts in Martínez-Sansigre et al. (2005). The error in flux (and luminosity) is the sum in quadrature of the photometric error (typically 10 per cent), the error from the slit (typically 5 per cent) and the uncertainty in the continuum. The FWHM is deconvolved by subtracting in quadrature the seeing from the observed size of the emission line. For lines with errors >20 per cent the measurements of the FWHM are not reliable, and the presence of broad lines (e.g. Mg II in AMS18) is therefore not significant.

Name	z	Line	λ_{rest} (Å)	λ_{obs} (Å)	FWHM (km s $^{-1}$)	Flux (W m $^{-2}$)	$\log_{10} L_{\text{line}}$ (W)	Error in flux (per cent)
AMS03	2.698 ± 0.009	Ly α	1216	4481	1400	1.22×10^{-19}	35.86	20
		Ly α	1216	4512	550	3.69×10^{-20}	35.34	18
AMS04	1.782 ± 0.001	Ly α	1216	3382	1800	5.01×10^{-19}	36.04	12
		[C III]	1909	5311	1450	1.02×10^{-19}	35.35	21
		[N II]	2142	5963	2550	1.16×10^{-19}	35.40	20
		[C II]	2326	6467	1250	8.59×10^{-20}	35.27	28
AMS05	2.017 ± 0.006	Ly α	1216	3658	1250	1.32×10^{-19}	35.59	38
		C IV	1549	4681	1000	1.16×10^{-19}	35.53	18
		He II	1640	4952	500	3.16×10^{-20}	34.97	41
AMS08	1.979 ± 0.004	Ly α	1216	3622	1900	5.05×10^{-19}	36.15	15
		C IV	1549	4622	1400	4.49×10^{-20}	35.10	23
		He II	1640	4877	900	2.30×10^{-20}	34.81	35
AMS12	2.767 ± 0.005	Ly α	1216	4586	650	4.41×10^{-19}	36.45	17
		C IV	1549	5824	1150	3.50×10^{-20}	35.35	25
		He II	1640	6181	1100	2.22×10^{-20}	35.15	26
AMS13	1.974 ± 0.003	Ly α	1216	3613	1650	9.88×10^{-19}	36.44	12
		C IV	1549	4611	1300	1.67×10^{-19}	35.67	20
AMS14	1.794 ± 0.001	Ly α	1216	3398	1500	3.69×10^{-19}	35.91	18
		C IV	1549	4326	600	5.61×10^{-20}	34.92	21
AMS16	4.169 ± 0.002	Ly α	1216	6288	1050	3.19×10^{-19}	36.73	12
		N V	1240	6406	2050	1.56×10^{-19}	36.42	28
		C IV	1549	8009	2050	2.69×10^{-19}	36.66	16
AMS17	3.137 ± 0.002	Ly α	1216	5031	1200	2.89×10^{-19}	36.40	16
AMS18	1.017 ± 0.004	[C II]	2326	4680	1400	8.19×10^{-20}	34.65	25
		Mg II	2798	5652	3500	1.14×10^{-19}	34.79	44
		[O II]	3727	7527	2200	1.12×10^{-19}	34.79	27

6 CRUDE PHOTOMETRIC REDSHIFTS

As discussed in Section 3, for $A_V \gtrsim 10$, the 3.6- μm flux density should be dominated by light from the host galaxy's old stellar population. We estimated crude 'photometric' redshifts for the entire sample by assuming the host galaxies to be the progenitors of present day massive elliptical galaxies. These galaxies were taken from the models of Bruzual & Charlot (2003), formed at $z = 10$ in a single starburst, with a Salpeter IMF (Salpeter 1955) and a stellar mass of $3.8 \times 10^{11} M_{\odot}$. Such host galaxies would become $2.6 L^*$ galaxies in the local K -band luminosity function (Cole et al. 2001). The justification for such an assumption is that radio-loud galaxies show a K - z Hubble diagram consistent with such massive host galaxies ($\sim 3L^*$; Jarvis et al. (2001b), Willott et al. (2003b)) and evidence points to the host galaxies of $z \sim 2$ radio-quiet quasars being $\sim 2L^*$ ellipticals (Kukula et al. 2001).

The scatter in true host luminosity means that individually the photometric redshifts are not necessarily close to the spectroscopic

redshift, but they agree on average for 10 of the 12 objects. There are two cases (AMS14 and AMS18) in which the photometric redshift is clearly overestimated. There are two possible explanations for this: a host galaxy which is smaller than the progenitors of ~ 2 – $3L^*$ ellipticals, and so our mass is an overestimate, or a star-forming host galaxy, in which case our elliptical galaxy template is not appropriate. For the objects that have spectroscopy, the mean and median values of z_{phot} are 2.75 and 2.65, respectively, while the mean and median spectroscopic redshifts are 2.33 and 2.00 (2.29 and 2.00 if the *Spitzer*–IRS are included); this represents reasonable agreement for such a simple photometric redshift estimation and gives us some confidence that we can estimate how many objects that remained unobserved or had blank optical spectra are at $z \geq 2$. To bring the median z_{phot} to the value of the median z_{spec} , one would have to change the mass of the elliptical to $2.9 \times 10^{11} M_{\odot}$, which would correspond to the progenitor of a $\sim 2L^*$ elliptical galaxy instead of $\sim 2.6L^*$. This would also bring the mean z_{phot} close to the mean




Figure 7. The spectrum of the type 1 quasar observed at the same time as AMS16.




Figure 8. Comparison between crude photometric and spectroscopic redshift for the objects with redshifts. Triangles are objects with narrow-lines in our spectroscopy; squares are objects with redshifts from *Spitzer*–IRS only. We see a large scatter and individually the photometric redshifts are of no use. On the whole, the agreement is reasonable except for two objects (AMS14 and AMS18). This means that the objects with spectroscopic redshifts in our sample are consistent with having massive elliptical galaxies as hosts.

z_{spec} . The fact that, for the objects with emission lines, the ‘photometric’ redshifts are roughly in agreement with the spectroscopic ones, suggests the host galaxies are the progenitors of massive ellipticals ($\gtrsim 2L^*$).

A possible source of error for our photometric redshifts would be the contribution of quasar light to $S_{3.6\mu\text{m}}$, together with a large host galaxy. This could plausibly make us miss some $z \sim 2$ type 2 quasars, which would only make the results of Martínez-Sansigre et al. (2005) more striking. Another source of error is if the host galaxy is not the progenitor of an elliptical, but a starburst (this possibility is explored in Section 9). The old stellar population would then be less luminous, and z_{phot} would be overestimated. The lack of lower redshift contaminants in our spectra gives us confidence that this is not a serious problem.

7 COMPOSITE TYPE 2 SPECTRUM

Following Rawlings, Eales & Lacy (2001), we proceeded to create a composite type 2 quasar spectrum by shifting the individual spectra to the rest frame with the IRAF task NEWREDSHIFT. The individual spectra were all scaled by their $\text{Ly}\alpha$ line flux and then averaged bin-by-bin with a sigma-clipping rejection algorithm with the clip

set to $\pm 1.5\sigma$. The object at $z = 1.02$ was therefore not included as there was no way of scaling it consistently. The composite of nine objects is shown in Fig. 9 and shows strong C IV (1549 Å) and He II (1640 Å) as well as $\text{Ly}\alpha$ (1216 Å) lines. [C II] (2326 Å) and Mg II (2798 Å) are also detected.

Table 4 shows the line fluxes relative to $\text{Ly}\alpha$. We can see C IV, He II and [C II] are clearly detected, with relative values of 0.36, 0.09 and 0.09, respectively. In addition, we see that the Mg II line is present, with a relative line flux of 0.05, while the N V (1240 Å) and [C III] (1909 Å) are not detected. The Mg II line must be detected at low SNR in several individual spectra to make it detected in the composite. [C III] is only detected in one of the objects making the composite and this could be due to the fact that at $z \gtrsim 2$ it is shifted to $\lambda \gtrsim 5700$ Å, where it falls near the dichroic cross-over region used in the 2004 July observations. AMS04 has [C III] and is at $z = 1.78$, but it was observed using a 5300-Å dichroic. The N V line is only present in AMS16. This composite spectrum is similar to that found for high-redshift radio galaxies (e.g. McCarthy 1993, and more recently Jarvis et al. 2001a; Rawlings et al. 2001), except that the relative strength of C IV is a factor of ~ 3 higher here than in McCarthy (1993). The [C II] line is stronger by a factor of 3 compared to McCarthy (1993) and the Jarvis et al. (2001a) value for radio galaxies with a projected linear size, $D > 70$ kpc, but consistent with the Jarvis et al. (2001a) value for radio galaxies with $D < 70$ kpc.

Comparing these ratios with McCarthy (1993), Rawlings et al. (2001) and Jarvis et al. (2001a), we find that they are broadly similar to those of powerful radio galaxies except C IV is significantly stronger, even stronger than for the McCarthy (1993) Seyfert 2 galaxies. The [C II] is similar to the Jarvis et al. (2001a) value for radio galaxies with $D < 70$ kpc. This might be an indication of a more complicated line emission mechanism than simple photoionization from a central engine, some of the ionization might be happening as a result of jet activity (see for example Best, Röttgering & Longair 2000; Jarvis et al. 2001a).

Comparing our composite type 2 to four individual high-redshift type 2s from the literature, we find that the Stern et al. (2002) type 2 ($z = 3.288$) has $\text{Ly}\alpha$, C IV and He II ratios similar to our composite (100:19:9), with slightly stronger C IV (relative to the other two lines). The Mainieri et al. (2005) object ($z = 3.660$) has a $\text{Ly}\alpha$ -to-C IV ratio (100:14) very similar to that of our composite and the Jarvis, van Breukelen & Wilman (2005) type 2 ($z = 1.65$) has a C IV-to-He II ratio (1:1) also in very good agreement with our composite. The Norman et al. (2002) type 2 at $z = 3.700$ has substantially weaker $\text{Ly}\alpha$ relative to C IV and He II (100:60:17); it is more similar to AMS16, for example, than to the composite. We therefore find very good agreement with the individual high-redshift type 2s from the literature and our sample.

Somewhat surprisingly another characteristic in common with radio galaxies is that all objects which show $\text{Ly}\alpha$ have at least some extended emission (in particular AMS03, AMS04, AMS08, AMS13 and AMS16 have very extended $\text{Ly}\alpha$ emission). AMS12 seems to have two $\text{Ly}\alpha$ components: a bright, spatially unresolved ‘peak’ sitting on top of a spatially extended fainter ‘plateau’. Also, AMS13 and AMS16 have other lines which are spatially extended (C IV and N V, respectively). In AMS04, the $\text{Ly}\alpha$ is ~ 5.5 arcsec in size (once deconvolved from the seeing), which is larger than the VLA beam size. For the other cases, we cannot currently tell whether the emission lines extended over regions larger than the radio emission. These characteristics all remind us of classic radio-loud sources despite their relatively low radio flux densities, so in Section 8 we proceed to compare our type 2 quasars with radio-loud objects.




Figure 9. Composite type 2 quasar spectrum. This has been made as described in Section 7, and it shows Ly α , particularly strong C IV, He II, [C II] and Mg II. [C III] is present in one individual object, and might be present in more but at $z \sim 2$ it falls around the dichroic cross-over region, so the noise might be too high to detect it. All the lines detected in the composite and in individual objects, except for Mg II, are present in the observations of IRAS FSC 10214+4724 (Serjeant et al. 1998).

Table 4. Comparison between the line ratios of this sample and those found by McCarthy (1993) for radio galaxies (RG) Seyfert 2s (SyII) and QSOs. The line fluxes are quoted as a percentage of the Ly α flux. The He II, C II and Mg II are roughly consistent with the radio galaxy values, but the C IV is significantly brighter, even brighter than found in Seyfert 2s.

Line	This sample	RG	McCarthy (1993) SyII	QSO
Ly α	100	100	100	100
C IV	36	12	22	8
He II	9	10	4	5
C II	9	3	–	4
Mg II	5	3	4	23

8 COMPARISON TO RADIO-LOUD OBJECTS

The mean radio flux density of the sample at 1.4 GHz is 780 μ Jy, which at the mean spectroscopic redshift $z = 2.33$ corresponds to a radio luminosity of $L_{1.4\text{GHz}} = 2 \times 10^{24} \text{ W Hz}^{-1} \text{ sr}^{-1}$ (assuming a spectral index $\alpha = 0.8$, where $L \propto \nu^{-\alpha}$). This is well below the break in the radio luminosity function (Willott et al. 1998) and in the regime where typical radio-selected objects have Fanaroff–Riley class I (FR I) radio structures (Fanaroff & Riley 1974). There is also little direct evidence of associated optical quasar activity (Grimes, Rawlings & Willott 2004) and weak or absent emission lines suggesting that, whether or not obscuring tori exist in this class of objects (Chiaberge, Capetti & Celotti 1999), any associated quasar is accreting at a very low rate (Rawlings & Saunders 1991). However, considering that FR Is have occasionally been found to accrete at high rates (Blundell & Rawlings 2001), we proceed to investigate the possibility of our objects having similar properties to radio-loud objects.

One typical characteristic of radio-loud objects is the presence of extended jets on scales larger than the host galaxy. Fig. 5 shows that most of the type 2 quasars shown in this paper are point-like in the VLA observations (~ 5 -arcsec resolution). In the few cases where the radio flux is extended, it generally coincides with another galaxy in the R band or 3.6- μ m images. However, if there is diffuse flux on large scales, the relatively long baselines of the VLA B-array could miss it out, so we would not see large jets.

The WSRT has also observed at 1.4 GHz the central region of the FLS, with an angular resolution of ~ 14 arcsec (Morganti et al. 2004). The shortest baselines of the WSRT are smaller than those of the VLA in B-array and this enables it to pick up flux on more extended scales, so any extended sources will have a significantly higher flux density in the WSRT catalogue than the VLA one. 13 of our targets are within the area observed by the WSRT: the fluxes are shown in Table 5. The rest of our sample was outside the area covered by the survey of Morganti et al. (2004). Most of the objects in Table 5 have consistent fluxes and therefore are unlikely to have a significant fraction of their radio flux in extended jets. AMS04, however, has a flux density as measured by WSRT which is 65 per cent higher than that measured by the VLA, suggesting the presence of jets. The 3.6- μ m radio overlay (Fig. 5) shows there is another radio source within the WSRT beam, and there is a hint of a faint galaxy which does not match the radio position perfectly. The companion radio source has a flux density of 331 μ Jy, so the WSRT flux density is clearly the sum of these two individual sources. If this is indeed due to another galaxy, then the WSRT is confusing both sources. However, we cannot be certain that the adjacent radio source is independent and due to another galaxy, so there is a chance that AMS04 has an extended jet with a flux density comparable to that of the central source.

The type 2s with the highest flux densities (AMS03, AMS10 and AMS13) all have more flux as observed by WSRT. Once again, however, there is a radio source with a clear optical (and infrared) ID next to one of the objects (AMS03), with a flux density of 203 μ Jy, so the WSRT flux is consistent with the sum of the two confused sources, within their errors. The flux of AMS13 is consistent in both telescopes to within 2σ and AMS10 is consistent within 3σ . Although 3σ might appear poor agreement, the reader is reminded of the highly spatially correlated nature of the noise in interferometer images, meaning the usual interpretation that only 0.5 per cent of measurements should differ by 3σ is not appropriate here. The central frequencies of both observations are not exactly the same: however, this small difference in frequency can only account for ~ 2 per cent of the flux difference.

To analyse the differences in flux density between both interferometers, we computed the normalized flux density difference: the measured flux density difference divided by the sum in quadrature of the 1σ errors of Table 5 ($(S_{\text{WSRT}} - S_{\text{VLA}})/(\sigma_{\text{WSRT}}^2 + \sigma_{\text{VLA}}^2)^{1/2}$, and plotted it in Fig. 10. For AMS03 and AMS04, we remove from the WSRT flux density the VLA flux density of the adjacent radio sources. Overplotted on the distribution are two Gaussian distributions, one with $\sigma = 1$, another with $\sigma = 2$. If there is no systematic difference between the two measurements, the plot should look like the Gaussian distribution with $\sigma = 1$, centred around 0. We see that this is clearly not the case, so the flux densities from both surveys are not consistent with random noise of variance $\sigma^2 = \sigma_{\text{WSRT}}^2 + \sigma_{\text{VLA}}^2$. However, the distribution roughly agrees somewhere between the $\sigma = 1$ and $\sigma = 2$ Gaussians (with $\sigma^2 = 2-4 \times (\sigma_{\text{WSRT}}^2 + \sigma_{\text{VLA}}^2)$). This suggests that the measurements from the WSRT and VLA are consistent provided the effective errors are taken to be larger (but still less than twice) the rms value of the noise. We therefore conclude that there is no firm evidence of extended radio emission around any of the 13 objects observed by both interferometers (other than possibly AMS04). The angular resolution of the VLA observations (5 arcsec) gives us an upper limit on the size of any jets. At $z = 2$, this corresponds to a jet size $\lesssim 40$ kpc.

Another known characteristic of radio-loud galaxies is the presence of extended line emission on similar scales to those of the radio jets. We have not only found that the composite spectrum resembles

Table 5. Comparison between the radio flux densities of the VLA in B-array (beam size ~ 5 arcsec) and the WSRT fluxes (beam size ~ 14 arcsec). We can see that the two flux densities agree within 3σ in all but two cases. Only AMS03 and AMS04 are not consistent, but examination of the VLA B-array images shows this is due to confusion. In the case of AMS03, it is clearly due to another radio source, while in the case of AMS04, it is not yet clear whether the confused flux density is from another source or from a diffuse radio component.

Name	$S_{\text{VLA } 1.4 \text{ GHz}}$ (μJy)	σ_{VLA} (μJy)	$S_{\text{WSRT } 1.4 \text{ GHz}}$ (μJy)	σ_{WSRT} (μJy)
AMS01	490	31	499	19
AMS02	1184	55	1209	17
AMS03	1986	87	2370	16
AMS04	536	72	887	13
AMS05	1038	49	1017	12
AMS06	444	31	380	14
AMS07	354	27	289	10
AMS08	655	36	679	12
AMS10	1645	73	1911	11
AMS11	356	29	253	9
AMS13	1888	83	2076	14
AMS18	390	29	351	23
AMS19	822	41	783	15




Figure 10. Normalized flux density difference distribution. This is computed as described in the test, and should match the Gaussian distribution with $\sigma = 1$ (inner Gaussian). This is clearly not the case, but it matches reasonably well somewhere between the $\sigma = 1$ and $\sigma = 2$ Gaussians.

that of radio galaxies, but several of our objects show significantly extended Ly α (AMS03, AMS04, AMS08, AMS13 and AMS16) and some have other extended lines (AMS13 and AMS16). In some objects (e.g. AMS16), the slit was placed along the direction of the slightly elongated radio axes, meaning that the emission lines could be extended along a hypothetical jet. In most cases, however, the slit PA was chosen for other reasons, so that any alignment with hypothetical jets would be coincidence. This suggests that any extended line emission is more likely to be associated with the remnants of a merger, which could have triggered the central engine, rather than an extended radio jet (something that has also been seen in radio galaxies, e.g., Villar-Martín et al. 2005).

The line ratios and the spatial extent of high-excitation lines of our objects seem to have some similarities to radio galaxies, as we have

discussed in Section 7. The [O III] line is an excellent indicator of the strength of the underlying quasar continuum (Simpson 1998a), so we now use the McCarthy (1993) radio galaxy line ratios to convert our line luminosities to [O III] (5007 Å) luminosities, to compare our results to the radio-loud samples discussed by Grimes et al. (2004). We converted the Ly α line to [O III], and when C IV is present we also converted that line. This yields two different estimates for the [O III] luminosity which are likely to be quite different, giving us a handle on the possible systematic errors in converting to [O III] luminosity, given that our composite spectrum has some differences with the McCarthy (1993) radio galaxy composite. The McCarthy (1993) ratios are $([\text{O III}]/\text{C IV})_{\text{RG}} = 2.38$ and $([\text{O III}]/\text{Ly}\alpha)_{\text{RG}} = 0.28$. We also convert radio flux at 1.4 GHz to luminosity at 151 MHz, assuming a spectral index $\alpha_{151}^{1.4} = 0.8$ (where flux density $\propto \nu^{-\alpha}$). The values obtained are shown in Table 6.

Fig. 11 shows how our objects sit on the L_{151} – $L_{[\text{O III}]}$ plane. The solid line shows the best-fitting line for the radio-loud galaxies from the 3CRR, 6CE and 7CRS samples (fig. 4 of Grimes et al. 2004). The region between the dashed lines encompasses all of the radio-loud objects, so all our objects except AMS03 lie in a different region to them. However, some important details must be remembered. First, we have assumed all our sources to be steep spectrum ($\alpha_{151}^{1400} = 0.8$), something for we have no current evidence. The true value of $\alpha_{151}^{1.4}$ is unlikely to be much larger (i.e. steeper) but objects with flatter true spectra will actually have lower L_{151} than shown here. In addition, the $L_{[\text{O III}]}$ for all of the objects has been estimated using the radio-galaxy ratios from McCarthy (1993) which have Ly α relative to C IV or He II systematically brighter than in our sample. The objects for which we have estimated $L_{[\text{O III}]}$ from Ly α (such as AMS03) are likely to have true $L_{[\text{O III}]}$ values brighter than those shown here. Finally, the dashed lines in Fig. 11 are not $\pm 1\sigma$ loci, they are the lines that encompass the entire radio-loud population (i.e. they could be considered $\sim 3\sigma$ contours). Once these details are taken into consideration, one sees that all the possible systematic errors would only lead to brighter [O III] lines and fainter luminosity at 151 MHz. Only two objects are anywhere near the radio-loud samples: AMS03 and AMS05. The first object has two different Ly α lines, and although we have used both for comparative reasons, the weaker of the two lines (the narrower line of the two) is unlikely to originate from an AGN, the stronger line should be looked at. The Ly α line of AMS05 has a large uncertainty in its flux measurement and is likely to be missing some flux since it was observed at a high airmass.

The sample presented here has typical [O III] luminosity (in logs) of 35–36. This is comparable to the radio-loud quasars in fig. 4 of Grimes et al. (2004), however, for a similar value of $L_{[\text{O III}]}$, the sample presented here has a L_{151} fainter by a factor of 100. It therefore differs from typical radio-selected objects in several respects: once the obscuration of the quasar nucleus is accounted for, they have the high accretion rates of typical quasars, but relatively low radio luminosities. The radio luminosities of these objects show that these are the radio-bright end of the radio-quiet population, or so called radio-intermediate quasars.

9 DISCUSSION

We have found a set of selection criteria that yields a population of radio-quiet type 2 quasars at $z \sim 2$. About half of the objects in this sample show emission lines in their optical spectra. The ratios of these emission lines are similar to those of radio galaxies with the main difference that the C IV line of type 2 quasars is much stronger and consistent with individual high-redshift type 2 quasars from the

Table 6. Data used to plot Fig. 11. The second column states which lines were used to estimate the [O III] line strength. The 151-MHz luminosities were estimated from the 1.4-GHz flux densities, assuming a spectral index $\alpha_{151}^{1.4} = 0.8$ between 151 MHz and 1.4 GHz.

Name	Line used	$\log_{10} L_{[\text{O III}]}$ (W)	$\log_{10} L_{151}$ (W Hz ⁻¹ sr ⁻¹)
AMS03	Ly α	35.3	25.6
AMS03	Ly α	34.8	25.6
AMS04	Ly α	35.5	24.7
AMS05	Ly α	35.0	25.1
AMS05	C IV	35.9	25.1
AMS08	Ly α	35.6	24.8
AMS08	C IV	35.5	24.8
AMS12	Ly α	35.9	25.3
AMS12	C IV	35.7	25.3
AMS13	Ly α	35.9	25.3
AMS13	C IV	36.0	25.3
AMS14	Ly α	35.4	24.4
AMS14	C IV	35.3	24.4
AMS16	Ly α	36.2	25.4
AMS16	C IV	37.0	25.4
AMS17	Ly α	35.8	25.3




Figure 11. Comparison to fig. 4 of Grimes et al. (2004). The symbols represent which line has been used to convert to [O III]: filled circles for Ly α , empty circles for C IV. The solid line represents the best fit to the radio-loud samples (3CRR, 6CE and 7CRS) in Grimes et al. (2004), with the dashed lines representing the loci which encompass all the radio-loud sources (so similar to $\pm 3\sigma$ contours).

literature. The Ly α is often spatially extended, as are sometimes the high ionization lines. This suggests that some of the line emission originates outside the classic narrow-line region, but whether this is due to jet activity or remnants of a merger cannot be deduced here. To further compare the type 2s to radio galaxies and radio-loud quasars, we compare their location on the L_{151} – $L_{[\text{O III}]}$ plane, and find significant differences. The type 2s have $L_{[\text{O III}]}$ comparable to radio-loud type 1 quasars, while being fainter at radio wavelengths by a factor of ~ 100 . In addition, the type 2s presented here show no signs of large extended jets (except possibly AMS04). Finally, comparison between our rough photometric redshift estimation, which assumes $2.6L^*$ host galaxies and our spectroscopic redshifts suggests that the objects in our sample which yield optical spectra are

consistent with having host galaxies which are massive ellipticals. In particular, changing our model elliptical from $2.6L^*$ to $2L^*$ would bring agreement to the median z_{phot} and z_{spec} , and close to agreement for the mean z_{phot} and z_{spec} . We have therefore found a population of classic (radio intermediate) type 2 quasars, showing only narrow lines with, presumably the torus blocking direct view of the quasar along our line of sight. We refer to these objects as ‘torus-obscured’ type 2s.

The lack of objects with spectroscopic redshifts around $z \sim 1.5$ could plausibly be due to the Ly α line being below the atmospheric cut-off, but C IV is detected in most of the higher z objects (and He II and [C II] are sometimes detectable) so the lack of objects at this redshift may instead be associated with the silicate absorption feature falling on the 24- μm band, as discussed in Section 2.

Of the remaining 11 objects, 10 show completely blank spectra and only one shows faint red continuum (AMS20). Their faint 3.6- μm flux densities and large R -band magnitudes suggest these are high redshift objects just like the ‘torus-obscured’ objects. In our 30-min integrations, type 1 quasars would have shown extremely bright continuum and lower redshift ($z \lesssim 1$) starbursts would have shown [O II] line emission or at least some strong continuum. Thus our sample does not suffer from any contamination: these blank objects are probably also high-redshift type 2 quasars.

Some of these 11 objects could be at $1.4 \leq z \leq 1.7$ with the C IV or He II lines too faint to be detected, although we have seen that 24- μm selection might disfavour this particular range of redshifts. The remaining objects are presumably obscured by dust on a large scale (~ 10 pc–1 kpc), which hides the narrow-line region as well as the broad-line region. Such concentration of dust is characteristic of star-forming regions, so we are presumably seeing either a nuclear or a galaxy-scale starburst. From their blank optical spectra, we know that these objects are indeed at high redshift and therefore their mid-infrared flux density is characteristic of objects accreting at a high rate, while their radio flux density is too high to originate only in star-forming regions and must be due to an AGN. In two cases (AMS06 and AMS19), published mid-infrared spectra confirm these objects to lie in the correct redshift range. In the other nine cases, despite the lack of lines to pinpoint their exact redshift, we are confident that these blank objects are type 2 quasars at high redshift. The obscuration, however, is not entirely due to an orientation effect, like for ‘torus-obscured’ objects, but due to the host galaxy, and so we refer to these objects as ‘host-obscured’ type 2s. At this stage, however, we cannot differentiate between dust at scales of ~ 10 pc or ~ 1 kpc. Such objects, lacking AGN emission lines in the optical spectra, have been found at lower redshift in the samples of Lacy et al. (2004, 2005a) and Leipski et al. (2005).

It is therefore not surprising that the Ly α (whether from the narrow-line or star-forming regions) in the host-obscured type 2s is too obscured for our exposure times (30 min) with a 4-m telescope: successful spectroscopy of submm galaxies in the blue requires very long integrations (1.5–6 h) with 8- or 10-m class telescopes (e.g. Chapman et al. 2005). We do note that AMS03 shows spatially extended Ly α , no other lines and no continuum, so there is a chance this object is host obscured and the second Ly α line traces the starburst: this requires further study.

In Martínez-Sansigre et al. (2005), the type 2 to type 1 ratio at $z \geq 2.00$ was modelled by predicting the number of type 1 quasars which would follow the 24- μm and radio criteria. The 3.6- μm criterion was simply assumed as a way of rejecting type 1s as well as providing a rough photometric redshift. The number of $z \geq 2.00$ type 1 quasars meeting the 24- μm was predicted from the B -band luminosity function of Wolf et al. (2003), converting from B band to 24 μm using




Figure 12. Schematic representation of the three types of quasar. The empty ellipses represent a transparent host galaxy, while the filled ellipses represent a dusty one. The solid circle represents the torus around the quasar. A is hosted by a transparent galaxy, and since we are viewing the torus face on, we can see the broad-line region. It is a type 1 quasar. B is also hosted by a transparent galaxy, but since the torus is edge on, we only see the narrow-line region. This is a torus-obscured type 2. C has a favourable geometry, while in D the torus is in the line of sight, but since both are obscured by large-scale dust in the galaxy, we cannot see the central region, so they are both host-obscured type 2s. From (Martínez-Sansigre et al. 2005), the ratio of B to A is $\sim 1:1$, while the ratio of B, C and D (together) to A is $\sim 2-3:1$. We note that although we are calling C and D ‘host’ obscured, the dust could plausibly be due to a nuclear starburst, with a characteristic scale ~ 100 pc, as well as a galaxy-scale starburst.

the Rowan-Robinson (1995) SED. The optical-to-radio correlation of Cirasuolo et al. (2003) was then used to predict what fraction of the $24\text{-}\mu\text{m}$ detected type 1s would also meet the radio criterion. This modelling predicted $4.3^{+2.2}_{-1.1}$ such type 1 quasars, at $z \geq 2.00$ and in our 3.8 deg^2 field. There are five narrow-line objects in our sample with spectroscopically confirmed $z \geq 2.00$, and the quasar type 2 to type 1 ratio for narrow-line (torus obscured) type 2s and type 1 quasars is therefore $\sim 1:1$. For the blank objects, the crude photometric redshifts can be used to estimate, on average, how many blank (host obscured) objects are at $z \geq 2$. There are six such objects, and so in (Martínez-Sansigre et al. 2005) the ratio of total number of type 2s (host and torus obscured) to type 1s was estimated to be $11/4$ or $\sim 3:1$. Since then AMS06 has been shown spectroscopically to lie at $z \leq 2.00$ the current number of objects at $z \geq 2.00$ is 10. From the errors in the modelled number of type 1 quasars, and the Poisson errors in our sample, the ratio of type 2 to type 1 quasars is found to be $2.3^{+1.8}_{-1.2}$. In a future paper, we will calculate better photometric redshifts, to try and obtain a more reliable estimate of the quasar fraction at $z \geq 2$.

The ‘two types of type 2’ quasar, and a type 1 quasar are schematically represented in Fig. 12. For a quasar to be of type 1 it has to lie in a relatively dust free galaxy (a ‘transparent’ galaxy) as well as having a favourable orientation to the torus (case A in Fig. 12). A quasar in a transparent galaxy, with an unfavourable angle to the torus will not outshine its host galaxy, but the narrow-line region will still be visible (case B: a torus-obscured type 2). When the host galaxy is dusty enough to obscure the quasar (cases C and D) it is no longer possible to see the central region, whether we have a favourable angle to the torus (C) or not (D). Objects of case C and D are likely to yield blank spectra, and are indistinguishable from each other with the current data set. Observations at radio frequencies, however, could shed some light on the orientation if radio jets are present.

This simple picture is likely to be applicable only to radio-quiet objects since the powerful radio jets are likely to be capable of removing dust on large scales (Baker et al. 2002; Willott et al. 2002). Therefore, although the ratio of type 2 to type 1 is likely to be $\sim 2-3:1$ for radio-quiet samples (and for all non-radio-selected samples since radio-quiet objects will dominate), it should be closer to $1:1$ for radio-loud samples. Thus, radio-loud samples will have few ‘host-obscured’ quasars (cases C and D, although some are probably present in the sample of Willott, Rawlings & Blundell 2001) and the quasar fraction measured in these radio-loud samples will be dominated by the geometry of the torus. This picture is therefore consistent with the ratio of narrow- to broad-line quasars predicted by unified models and found for radio-loud objects ($\sim 1:1$ e.g. Willott et al. 2000).

Inferring a type 2 to type 1 ratio $\sim 2-3:1$ for the sample presented in this article does not disagree with the receding-torus models (e.g. Lawrence 1991; Simpson 2005), since the obscuring mechanism for ‘host-obscured’ type 2s is unrelated to the torus orientation or geometry. The ratio of torus-obscured (B) to type 1 (A) quasars predicted for our sample is $\sim 1:1$ (and indeed consistent with the receding torus: fig. 2 of Martínez-Sansigre et al. 2005). This phenomenon of host obscuration might also be present in the Seyfert 2s at $z \sim 1$ that dominate the hard X-ray background, in which case care must be taken to separate the obscuring mechanism necessary to explain the AGN dominating the X-ray background from modifications to the unified schemes such as the receding torus.

The ratio of $\sim 2-3:1$ found for this sample is similar to the ratio required for Seyfert 2s to Seyfert 1s at $z \sim 1$ to fit the hard X-ray background, although a ratio $\sim 2-3:1$ for the quasars is not required by the hard X-ray background (Ueda et al. 2003; Treister & Urry 2005). We do note that, due to the range in gas-to-dust ratios, some of the X-ray obscured quasars that contribute to the hard X-ray background are not necessarily type 2s in the optical sense (they might only be reddened quasars with $A_V \sim 1-3$ but a large gas-to-dust ratio). However, at least in our sample, we find the ratio of all type 2s (B, C and D) to type 1 (A) is $\sim 2-3:1$: a higher type 2 to type 1 ratio than is found in deep X-ray surveys. This suggests that indeed, $24\text{-}\mu\text{m}$ selection can find type 2 quasars that X-ray selection in the $2-10\text{ keV}$ band has not been able to detect, and therefore a number of the type 2 quasars presented here should be Compton thick. This, of course, can be tested via X-ray follow-up of this or a similar sample.

ACKNOWLEDGMENTS

We would like to warmly thank Richard Wilman for access to his absorbed X-ray spectra and for useful comments, the anonymous referee for valuable suggestions, and Caroline van Breukelen for help with the overlays. AMS would like to thank the Council of the European Union for support; SR and CS would like to thank the UK PPARC for a Senior Research Fellowship and an Advanced Fellowship, respectively. This work is based on observations made with the *Spitzer* Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology. The William Herschel Telescope (WHT) is operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatory del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

REFERENCES

- Alexander D. M. et al., 2003, *AJ*, 126, 539
- Alexander D. M., Chartas G., Bauer F. E., Brandt W. N., Simpson C., Vignali C., 2005, *MNRAS*, 357, L16

- Antonucci R., 1993, *ARA&A*, 31, 473
- Baker J. C., Hunstead R. W., Athreya R. M., Barthel P. D., de Silva E., Lehnert M. D., Saunders R. D. E., 2002, *ApJ*, 568, 592
- Best P. N., Röttgering H. J. A., Longair M. S., 2000, *MNRAS*, 311, 23
- Blundell K. M., Rawlings S., 2001, *ApJ*, 562, L5
- Bohlin R. C., Savage B. D., Drake J. F., 1978, *ApJ*, 224, 132
- Bouchet P., Lequeux J., Maurice E., Prevot L., Prevot-Burnichon M. L., 1985, *A&A*, 149, 330
- Brandl B. R. et al., 2004, *ApJS*, 154, 188
- Broadhurst T., Lehar J., 1995, *ApJ*, 450, L41
- Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
- Chapman S. C., Blain A. W., Smail I., Ivison R. J., 2005, *ApJ*, 622, 772
- Chiaberge M., Capetti A., Celotti A., 1999, *A&A*, 349, 77
- Cirasuolo M., Celotti A., Magliocchetti M., Danese L., 2003, *MNRAS*, 346, 447
- Cole S. et al., 2001, *MNRAS*, 326, 255
- Comastri A., Setti G., Zamorani G., Hasinger G., 1995, *A&A*, 296, 1
- Condon J. J., 1992, *ARA&A*, 30, 575
- Condon J. J., Cotton W. D., Yin Q. F., Shupe D. L., Storrie-Lombardi L. J., Helou G., Soifer B. T., Werner M. W., 2003, *AJ*, 125, 2411
- Croom S. M., Smith R. J., Boyle B. J., Shanks T., Miller L., Outram P. J., Loaring N. S., 2004, *MNRAS*, 349, 1397
- Czerny B., Li J., Loska Z., Szczerba R., 2004, *MNRAS*, 348, L54
- Fadda D., Jannuzi B. T., Ford A., Storrie-Lombardi L. J., 2004, *AJ*, 128, 1
- Fadda D. et al., 2006, preprint (astro-ph/0603488)
- Fanaroff B. L., Riley J. M., 1974, *MNRAS*, 167, 31p
- Gaskell C. M., Goosmann R. W., Antonucci R. R. J., Whysong D. H., 2004, *ApJ*, 616, 147
- Gilli R., Salvati M., Hasinger G., 2001, *A&A*, 366, 407
- Granato G. L., Danese L., 1994, *MNRAS*, 268, 235
- Grimes J. A., Rawlings S., Willott C. J., 2004, *MNRAS*, 349, 503
- Hopkins et al., 2004, *AJ*, 128, 1112
- Jarvis M. J., Rawlings S., Eales S., Blundell K. M., Bunker A. J., Croft S., McLure R. J., Willott C. J., 2001a, *MNRAS*, 326, 1563
- Jarvis M. J., Rawlings S., Eales S., Blundell K. M., Bunker A. J., Croft S., McLure R. J., Willott C. J., 2001b, *MNRAS*, 326, 1585
- Jarvis M. J., van Breukelen C., Wilman R. J., 2005, *MNRAS*, 358, L11
- Kukula M. J., Dunlop J. S., McLure R. J., Miller L., Percival W. J., Baum S. A., O'Dea C. P., 2001, *MNRAS*, 326, 1533
- Lacy M. et al., 2004, *ApJS*, 154, 166
- Lacy M., Canalizo G., Rawlings S., Sajina A., Storrie-Lombardi L., Armus L., Marleau F. R., Muzzin A., 2005a, *Mem. Soc. Astron. Ital.*, 76, 154
- Lacy M. et al., 2005b, *ApJS*, 161, 41
- Lawrence A., 1991, *MNRAS*, 252, 586
- Leipski C. et al., 2005, *A&A*, 440, L5
- Lonsdale C. J. et al., 2003, *PASP*, 115, 897
- Madau P., Ghisellini G., Fabian A. C., 1994, *MNRAS*, 270, L17
- Mainieri V. et al., 2005, *MNRAS*, 356, 1571
- Maiolino R., Marconi A., Salvati M., Risaliti G., Severgnini P., Oliva E., La Franca F., Vanzi L., 2001, *A&A*, 365, 28
- Marleau F. R. et al., 2004, *ApJS*, 154, 66
- Martínez-Sansigre A., Rawlings S., Lacy M., Fadda D., Marleau F. R., Simpson C., Willott C. J., Jarvis M. J., 2005, *Nat*, 436, 666
- McCarthy P. J., 1993, *ARA&A*, 31, 639
- Morganti R., Garrett M. A., Chapman S., Baan W., Helou G., Soifer T., 2004, *A&A*, 424, 371
- Norman C. et al., 2002, *ApJ*, 571, 218
- Pei Y. C., 1992, *ApJ*, 395, 130
- Rawlings S., Saunders R., 1991, *Nat*, 349, 138
- Rawlings S., Eales S., Warren S., 1990, *MNRAS*, 243, 14p
- Rawlings S., Eales S., Lacy M., 2001, *MNRAS*, 322, 523
- Richards G. T. et al., 2002, *AJ*, 123, 2945
- Richards G. T. et al., 2003, *AJ*, 126, 1131
- Rowan-Robinson M., 1995, *MNRAS*, 272, 737
- Rowan-Robinson M. et al., 1991, *Nat*, 351, 719
- Salpeter E. E., 1955, *ApJ*, 121, 161
- Sanders D. B., Mirabel I. F., 1996, *ARA&A*, 34, 749
- Serjeant S., Rawlings S., Lacy M., McMahon R. G., Lawrence A., Rowan-Robinson M., Mountain M., 1998, *MNRAS*, 298, 321
- Simpson C., 1998a, *MNRAS*, 297, L39
- Simpson C., 1998b, *ApJ*, 509, 653
- Simpson C., 2005, *MNRAS*, 360, 565
- Simpson C., Rawlings S., Lacy M., 1999, *MNRAS*, 306, 828
- Stern D. et al., 2002, *ApJ*, 568, 71
- Treister E., Urry C. M., 2005, *ApJ*, 630, 115
- Treister E. et al., 2004, *ApJ*, 616, 123
- Treister E. et al., 2006, *ApJ*, 640, 603
- Ueda Y. et al., 1999, *ApJ*, 518, 656
- Ueda Y., Akiyama M., Ohta K., Miyaji T., 2003, *ApJ*, 598, 886
- Urrutia T., Lacy M., Gregg M. D., Becker R. H., 2005, *ApJ*, 627, 75
- van Ojik R., Roettgering H. J. A., Miley G. K., Hunstead R. W., 1997, *A&A*, 317, 358
- Villar-Martín M., Tadhunter C., Morganti R., Holt J., 2005, *MNRAS*, 359, L5
- Weedman D. W., Le Floch E., Higdon S. J. U., Higdon J. L., Houck J. R., 2006, *ApJ*, 638, 613
- Werner M. W. et al., 2004, *ApJS*, 154, 1
- Wilkes B. J., Pounds K. A., Schmidt G. D., Smith P. S., Cutri R. M., Ghosh H., Nelson B., Hines D. C., 2005, *ApJ*, 634, 183
- Willott C. J., 2005, *ApJ*, 627, L101
- Willott C. J., Rawlings S., Blundell K. M., Lacy M., 1998, *MNRAS*, 300, 625
- Willott C. J., Rawlings S., Blundell K. M., Lacy M., 2000, *MNRAS*, 316, 449
- Willott C. J., Rawlings S., Blundell K. M., 2001, *MNRAS*, 324, 1
- Willott C. J., Rawlings S., Archibald E. N., Dunlop J. S., 2002, *MNRAS*, 331, 435
- Willott C. J. et al., 2003a, *MNRAS*, 339, 397
- Willott C. J., Rawlings S., Jarvis M. J., Blundell K. M., 2003b, *MNRAS*, 339, 173
- Willott C. J. et al., 2004, *ApJ*, 610, 140
- Wilman R. J., Fabian A. C., 1999, *MNRAS*, 309, 862
- Wolf C., Wisotzki L., Borch A., Dye S., Kleinheinrich M., Meisenheimer K., 2003, *A&A*, 408, 499
- Worsley M. A. et al., 2005, *MNRAS*, 357, 1281
- Yan L. et al., 2005, *ApJ*, 628, 604
- Zheng W. et al., 2004, *ApJS*, 155, 73

This paper has been typeset from a \LaTeX file prepared by the author.